NCH690

300120

GENERAL INFORMATION

Description

The NCH640 cell line is a glioblastoma stem-like cell model used in research to explore the mechanisms of tumor resistance, cellular survival under stress, and therapeutic responses. Glioblastoma, one of the most aggressive forms of brain tumors, is difficult to treat due to its resistance to therapy and adaptation to a hostile microenvironment. NCH640 is cultured in specialized media such as Neurobasal A with supplements like B27, and its growth is supported by essential growth factors like EGF and FGF-2. It is often used alongside other glioma stem cell models, such as NCH690 and NCH644, to investigate these biological phenomena.
The research on NCH640 focuses heavily on its resistance mechanisms, particularly under hypoxic conditions. Glioma cells like NCH640 show significant reliance on metabolic adaptations, including altered reactive oxygen species (ROS) regulation. Studies have demonstrated that targeting pathways such as the integrated stress response (ISR) in NCH640 and related cell lines may improve their sensitivity to therapies like temozolomide, which is commonly used in glioblastoma treatment. These findings are important for devising new strategies to overcome the inherent resistance of glioma stem-like cells to standard therapeutic interventions.

Tissue	Brain
Disease	Glioblastoma

CLASSIFICATION

Organism	Human
Sex	Female
Growth properties	Adherent
Biosafety level	1

CHARACTERISTICS

Tumourigenic	Yes

IDENTIFIERS

Citation

NCH690 (Cytion catalog number 300120)

NCBI TaxID	9606
Cellosaurus	CVCL_x915

HANDLING AND CULTURE

Culture medium

DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO3 (Cytion article number 820400a)

Incubation atmosphere	37°C, 5% CO_2, humidified atmosphere.

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Supplements

Supplement the medium with 10% FBS, 5 mg/L Heparin, 20 ng/mL bFGF, 20 microgram/L EGF, 5 mg/L Insulin, 100 mg/L Transferrin, 5,2 microgram/L Na-selenit, 6,3 microgram/L Progesteron, 161,1 microgram/L Putrescin, 50 mg/L Hydrocortinson

Subculturing

For subculturing spheroid cultures, begin by mechanically dissociating the spheroids through pipetting up and down 5 to 10 times using an Eppendorf pipette with 1000 µl filter tips. After this, centrifuge the mixture at 300g for 5 minutes at room temperature to pellet the cells. Discard the supernatant and resuspend the cell pellet in fresh culture medium. Finally, transfer the resuspended cells into new culture vessels to promote further spheroid formation. This approach ensures efficient spheroid breakdown and readies them for continued growth in a new environment.

Seeding density	1 x 10^5 cells/mL
Fluid renewal	2 to 3 times per week
Post-thaw recovery	After thawing allow the cells to recover from the freezing process for at least 24 to 48 hours.

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Freeze medium	As a cryopreservation medium, we use 50% basal medium + 40% FBS + 10% DMSO, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Storage conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Shipping conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

DONOR

Donor age	78 years
Ethnicity	Caucasian

LEGAL AND SAFETY

Safety precautions

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Warranty

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Third-party agreements apply	Yes

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB