CLS-439 □□

300150

□ CLS □ 1998 □□□□ 61 □□□□□□□□□□□□□□□□□

GENERAL INFORMATION

Description

□ CLS □ 1998 □□□□ 61 □□□□□□□□□□□□□□□□□

Tissue	□□
Disease	□

Synonyms

- CLS439

CLASSIFICATION

Organism	Human
Sex	Male
Growth properties	Adherent
Biosafety level	1

CHARACTERISTICS

Morphology	□□□
Tumourigenic	□□□□□□□□
Doubling time	35 □□

IDENTIFIERS

Citation

CLS-439□Cytion □□ 300150□

NCBI TaxID	9606
Cellosaurus	CVCL_5982

HANDLING AND CULTURE

Culture medium

□□□ 5a □□□□□□3.0 g/L □□□□□□□□□□□□□□□□□2.0 mM □□□□□□□2.2 g/L $NaHCO_3$□Cytion □□□□ 820200a□

Incubation atmosphere	37°C□5% CO_2□□□□□□

Supplements

□□□□□□□ 10% □□□□□□FBS□

Dissociation reagent	Accutase

Subculturing
移除培養基，並使用不含鈣和鎂的PBS沖洗附著細胞（T25細胞培養瓶需3-5毫升PBS，T75細胞培養瓶需5-10毫升PBS）。 加入 TrypleExpress（T25 培養瓶每瓶 1–2 毫升，T75 培養瓶每瓶 2.5 毫升），務必完全覆蓋細胞片。於室溫下孵育 10 分鐘。 小心地將細胞重新懸浮，可選擇性添加培養基（非必要），並分裝至裝有新鮮培養基的新培養瓶中。

Seeding density 1 × 10⁴ 個細胞/cm² 約需 3 天即可形成密合的細胞層

Fluid renewal 每週 2 至 3 次

Post-thaw recovery 細胞在 37 攝氏度／5% CO₂ 條件下解凍後，必須靜置至少 24 小時

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility
透過基於 PCR 的檢測及基於發光原理的支原體檢測方法，排除支原體污染的可能性。
為確保無細菌、真菌或酵母菌污染，每日均會對細胞培養物進行目視檢查。

Freeze medium 作為冷凍保存培養基，我們使用完全生長培養基（含 FBS）+ 10% DMSO 以確保解凍後的存活率，或使用 CM-1
（Cytion 型號 800100），該培養基含有經優化的滲透保護劑和代謝穩定劑，可提升細胞恢復率並減輕冷凍誘導的壓
力。

Storage conditions
若需長期保存，請將試管置於溫度約為 −150 至 −196 °C 的氣相液氮中。僅在轉移至液氮之前，可將試管暫時存放於 −80 °C 環境中，作為短期過渡步驟。

Shipping conditions
冷凍保存的細胞系會以乾冰冷藏，並裝於經驗證的隔熱包裝中，包裝內含有足夠的冷媒，以確保在運輸過程中溫度維持在約 −78 °C。收到貨物後，請立即檢查容器，並盡快將試管轉移至適當的儲存場所。

DONOR

Donor age 61年

Ethnicity 高加索人

LEGAL AND SAFETY

Safety precautions
若計劃將低溫試管存放在液態氮中以供日後解凍，必須嚴格遵守相關安全措施。 必須穿戴適當的防護手套和防護服，且在將冷凍樣本移入或移出液氮罐時，必須佩戴口罩或安全護目鏡。此舉旨在降低因取出時冷凍試管可能發生爆炸而導致受傷的風險，此類爆炸可能造成尖銳碎片飛濺。

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

2 / 3

Warranty

□ Cytion□□□□□□□□□□□□□□□□□□□□
□□□□□□□□□□□□□□□
60 □□□□□□□□□□□□□□□□□□□□□□□
□□□□□□□□□□□□
□□□□□□□□□□□□□□□□□□□□□
□□□□□□□□
□□□□□□□□

Material Transfer Agreement required	Yes