DAN-G

300162

GENERAL INFORMATION

Description

The DAN-G cell line is derived from a human pancreatic carcinoma. It is extensively utilized in research focused on pancreatic cancer, particularly in studies pertaining to tumorigenesis, metastasis, and chemotherapy resistance. The genetic profile of DAN-G includes mutations in key oncogenes and tumor suppressor genes, which are characteristic of pancreatic adenocarcinomas. This makes the cell line a valuable model for understanding the molecular mechanisms underlying pancreatic cancer and for testing new therapeutic strategies.
In addition to its applications in cancer research, the DAN-G cell line has been used to study the cellular processes involved in the progression of pancreatic ductal adenocarcinoma, including cell cycle regulation, apoptosis, and signal transduction pathways. The cells exhibit aggressive in vitro growth characteristics and have the ability to form tumors in immunocompromised mice, which simulates the human disease and provides an in vivo system for evaluating the efficacy of anticancer drugs. Researchers also employ this cell line to investigate the role of the tumor microenvironment in pancreatic cancer progression and resistance to therapy.

Tissue	Pancreas
Disease	Adenocarcinoma

Synonyms

- Dan-G
- DanG
- DANG

CLASSIFICATION

Organism	Human
Sex	Female
Growth properties	Adherent
Biosafety level	1

CHARACTERISTICS

Morphology	Epithelial-like
Tumourigenic	Yes, in nude mice
Doubling time	33 hours

IDENTIFIERS

Citation

DAN-G (Cytion catalog number 300162)

NCBI TaxID	9606
Cellosaurus	CVCL_0243

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

EXPRESSION AND SUSCEPTIBILITY

Protein expression

P53 negative

Mutational profile

DAN-G cells carry a homozygous Kras mutation in codon12: GGT(Gly) >GTT(Val)

HANDLING AND CULTURE

Culture medium

RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO3 (Cytion article number 820700a)

Incubation atmosphere	37°C, 5% CO_2, humidified atmosphere.

Supplements

Supplement the medium with 10% FBS

Dissociation reagent	Accutase

Subculturing

Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Seeding density	3 to 4 x 10^4 cells/cm^2 will yield in a confluent layer in about 4 days
Fluid renewal	2 to 3 times per week
Post-thaw recovery	After thawing, plate the cells at 5 x 10^4 cells/cm^2 and allow the cells to recover from the freezing process and to adhere for at least 24 hours.

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Storage conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Shipping conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

DONOR

Donor age	68 years

LEGAL AND SAFETY

Safety precautions

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Warranty

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Material Transfer Agreement required	Yes