U-2 OS-CRISPR-NUP96-mEGFP clone no.195

300174

GENERAL INFORMATION

Description

The U-2 OS-CRISPR-NUP96-mEGFP is a genetically modified cell line derived from the human osteosarcoma U-2 OS parent line. This cell line incorporates a targeted insertion of the monomeric Enhanced Green Fluorescent Protein (mEGFP) tag at the NUP96 gene locus, achieved through the CRISPR-Cas9 gene-editing technology. NUP96, part of the nuclear pore complex, is essential for nuclear transport, and its fusion with mEGFP allows for real-time visualization of nuclear pore dynamics under fluorescent microscopy, providing valuable insights into nuclear transport mechanisms and nucleocytoplasmic trafficking.
This specific clone, numbered 195, has been selected for its stable expression of the NUP96-mEGFP fusion protein and maintains the typical characteristics of the U-2 OS lineage, including a robust cytoskeletal structure which is critical in studies related to cancer cell migration and metastasis. The application of CRISPR technology ensures precise gene editing, minimizing off-target effects which could compromise the integrity of the experimental outcomes. This makes U-2 OS-CRISPR-NUP96-mEGFP clone no.195 particularly useful for high-resolution imaging techniques and detailed cellular architecture studies, aiding in advanced research in cellular biology, cancer research, and nuclear transport phenomena.

Tissue	Bone
Disease	Osteosarcoma

CLASSIFICATION

Organism	Human
Sex	Female
Growth properties	Adherent
GMO status	GMO-S1

GMO status details

GMO-S1: This human osteosarcoma cell line (U2OS-CRISPR-NUP96-mEGFP, clone 195) contains a CRISPR-engineered NUP96-mEGFP fusion introduced via lentiviral delivery, enabling fluorescent tracking of nuclear pore complexes. The modification is stably integrated. This classification applies only within Germany and may differ elsewhere.

Biosafety level	1

CHARACTERISTICS

Morphology	Epithelial-like

IDENTIFIERS

Citation

U-2 OS-CRISPR-NUP96-mEGFP clone no.195 (Cytion catalog number 300174)

NCBI TaxID	9606
Cellosaurus	CVCL_B7FJ
Depositor	The Ellenberg Lab (EMBL)

EXPRESSION AND SUSCEPTIBILITY

Protein expression

MEGFP (nuclear pore complex protein 96, mEGFP tagged)

HANDLING AND CULTURE

Culture medium

McCoys 5a, w: 3.0 g/L Glucose, w: 1.5 mM Glutamine, w: 2.2 g/L $NaHCO_3$ (Cytion article number 820200a)

Incubation atmosphere	37°C, 5% CO_2, humidified atmosphere.

Supplements

Supplement the medium with 10% FBS, 1% NEAA and 1.0 mM Sodium pyruvate

Dissociation reagent	Accutase

Subculturing

Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Seeding density	2 to 3 x 10^4 cells/cm^2
Fluid renewal	2 to 3 times per week

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Storage conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Shipping conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

DONOR

Donor age	15 years
Ethnicity	Caucasian

LEGAL AND SAFETY

Safety precautions

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Warranty

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Third-party agreements apply	Yes