MeWo

300285

GENERAL INFORMATION

Description

The MeWo cell line is a fibroblast-like melanoma cell line isolated from the skin of a 78-year-old White male patient with malignant melanoma. These cells exhibit a characteristic morphology that reflects their fibroblastic origin. MeWo cells are valuable in cancer research, particularly for studying melanoma's biological properties and immune interactions. As with other melanoma cell lines, MeWo cells have been instrumental in the study of tumor antigens and their immunogenicity. Various studies have utilized MeWo cells to identify specific surface antigens, which are crucial in understanding how melanoma cells interact with the immune system.

One of the notable properties of MeWo cells is their ability to support the growth of varicella-zoster virus (VZV) isolates, with optimal growth conditions at 32°C, although they can still sustain VZV growth at 36°C. This makes the MeWo cell line particularly useful in virological research, especially in the context of viral replication and pathogenesis studies under varying temperature conditions. Additionally, MeWo cells are tumorigenic, as they can form tumors when injected into nude mice, a property that underscores their utility in in vivo tumorigenicity studies. This characteristic, coupled with their responsiveness to viral infection, highlights MeWo cells as a versatile model for both cancer and infectious disease research.

Studies involving the MeWo cell line have also explored the expression of melanoma-associated antigens, where MeWo has been used as a reference cell line in absorption assays to identify unique and common antigens across different melanoma samples. The antigenic profile of MeWo cells, as identified in these studies, includes antigens that are shared with other melanoma cell lines, as well as those that may be unique to this cell line, contributing to the broader understanding of melanoma immunology.

Tissue	Skin
Disease	Cutaneous melanoma
Metastatic site	Lymph node

Applications

Virus studies

Synonyms

- MEWO
- Mewo
- Me Wo
- Me-Wo
- Mevo
- SK-MEL-MeWo
- Mel-MeWo
- BI-Mel
- EST50

CLASSIFICATION

Organism	Human
Sex	Male
Growth properties	Adherent
Biosafety level	1

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

CHARACTERISTICS

Morphology	Fibroblast-like
Tumourigenic	Forms malignant melanoma
MSI status	Stable (MSS)

IDENTIFIERS

Citation

MeWo (Cytion catalog number 300285)

NCBI TaxID	9606
Cellosaurus	CVCL_0445

EXPRESSION AND SUSCEPTIBILITY

Mutational profile

BRAF V600E wt

Products

Melanin

HANDLING AND CULTURE

Culture medium

EMEM (MEM Eagle), w: 2 mM L-Glutamine, w: 2.2 g/L NaHCO3, w: EBSS (Cytion article number 820100a)

Incubation atmosphere	37°C, 5% CO_2, humidified atmosphere.

Supplements

Supplement the medium with 10% FBS and 1% NEAA

Dissociation reagent	Accutase

Subculturing

Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Fluid renewal	2 to 3 times per week

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Storage conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Shipping conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

DONOR

Donor age	78 years
Ethnicity	Caucasian

LEGAL AND SAFETY

Safety precautions

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Warranty

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Material Transfer Agreement required	Yes