KMS-12-BM

300287

GENERAL INFORMATION

Description

The KMS-12-BM cell line is a human myeloma cell line established from the bone marrow of a patient with non-producing multiple myeloma. This cell line represents an immature plasmacytoid stage of B-cell differentiation, characterized by the expression of surface markers CD20, CD38, and PCA-1, but a lack of immunoglobulin production. The cells are notable for their distorted morphology, with many showing multinuclear and giant characteristics. Ultrastructurally, KMS-12-BM cells possess well-developed rough endoplasmic reticulum and ovoid eccentric nuclei with peripheral chromatin distribution, typical of plasmacytoid cells.
KMS-12-BM cells exhibit a chromosomal abnormality, particularly a reciprocal translocation t(11;14)(q13;q32), which is often associated with multiple myeloma. These cells also display a broad range of chromosomal numbers, from hypodiploid to polyploid, indicating significant genomic instability. Unlike its counterpart KMS-12-PE, the KMS-12-BM line does not produce amylase, and it lacks immunoglobulin secretion or surface expression, making it suitable for studies involving immunoglobulin-nonproducing myeloma. Additionally, it shows low cloning efficiency in soft agar culture conditions, with less than 0.1% colony formation, and has no tumorigenic properties when injected into nude mice.

Tissue	Bone marrow
Disease	Multiple Myeloma

Synonyms

- KMS 12 BM
- KMS-12BM
- KMS12-BM
- KMS12BM
- KMS-12
- KMS12
- Kawasaki Medical School-12-Bone Marrow

CLASSIFICATION

Organism	Human
Sex	Female
Growth properties	Suspension
Biosafety level	1

CHARACTERISTICS

Morphology	Round cells
Cell type	B cell
Tumourigenic	Not tumorigenic in nude mice

IDENTIFIERS

Citation

KMS-12-BM (Cytion catalog number 300287)

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

NCBI TaxID	9606
Cellosaurus	CVCL_1334

EXPRESSION AND SUSCEPTIBILITY

Surface antigens

CD3 -, CD10 -, CD13 -, CD19 -, CD20 +, CD34 -, CD37 +, CD38 +, cyCD79a +, CD80 -, CD138 +, HLA-DR -, PCA-1 +, sm/cyIgG -, sm/cyIgM -, sm/cykappa -, sm/cylambda -

Mutational profile

Translocation: t(11;14)(q13;q32)

Products

No immunoglobulin production

HANDLING AND CULTURE

Culture medium

RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO3 (Cytion article number 820700a)

Incubation atmosphere	37°C, 5% CO_2, humidified atmosphere.

Supplements

Supplement the medium with 10% FBS

Subculturing

Maintain cultures by periodically adding or replacing the medium. Initiate cultures with a density of 5 x 10^5 cells/ml and keep the cell concentration within the range of 3 x 10^5 to 1 x 10^6 cells/ml for optimal growth.

Seeding density	5 x 10^5 cells/ml

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Storage conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Shipping conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

DONOR

Donor age	64 years
Ethnicity	Japanese

LEGAL AND SAFETY

Safety precautions

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Warranty

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Material Transfer Agreement required	Yes