# B16-F0

300308

## GENERAL INFORMATION

### Description

The B16-F0 cell line is a murine melanoma cell line derived from the B16 mouse melanoma. This cell line is widely used in cancer research due to its high metastatic potential and ability to form tumors when injected into syngeneic mice. B16-F0 cells are particularly useful for studying the molecular mechanisms underlying melanoma progression and metastasis, as well as for testing the efficacy of anti-cancer drugs and therapeutic interventions in preclinical models. Notably, the B16-F0 cell line is the parent cell line from which other variants, such as B16-F1, B16-F10, and B16-BL6, have been derived through selective procedures aimed at enhancing specific metastatic properties.
B16-F0 cells exhibit a typical epithelial morphology and grow adherently in culture. They are known to express various melanoma-associated antigens, making them a valuable tool for immunological studies and for developing melanoma vaccines. Additionally, these cells are often employed in studies involving gene expression, signaling pathways, and the tumor microenvironment. Researchers utilize B16-F0 cells to explore the interactions between melanoma cells and the immune system, particularly focusing on immune evasion and suppression mechanisms. The characterization of B16-F0 and its derived lines provides a comprehensive framework for understanding the invasive and metastatic behaviors of melanoma, with B16-F1, B16-F10, and B16-BL6 each representing stages of increasing metastatic and invasive activity, thereby serving as critical models in the study of cancer progression and therapeutic response.

| | |
|---|---|
| **Tissue** | Skin |
| **Disease** | Mouse melanoma |

### Synonyms

- B16/F0
- B16F0

## CLASSIFICATION

| | |
|---|---|
| **Organism** | Mouse |
| **Sex** | Male |
| **Growth properties** | Adherent |
| **Biosafety level** | 1 |

## CHARACTERISTICS

| | |
|---|---|
| **Morphology** | Mixture of spindle-shaped and epithelial-like cells |
| **Cell type** | Epithelial |
| **Tumourigenic** | Yes, in syngeneic mice |

## IDENTIFIERS

### Citation

B16-F0 (Cytion catalog number 300308)

| | |
|---|---|
| **NCBI TaxID** | 10090 |
| **Cellosaurus** | CVCL_0604 |

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

## EXPRESSION AND SUSCEPTIBILITY

**Products**

Melanin

## HANDLING AND CULTURE

**Culture medium**

DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO3, w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)

| | |
|---|---|
| **Incubation atmosphere** | 37°C, 5% $CO_2$, humidified atmosphere. |

**Supplements**

Supplement the medium with 10% FBS

| | |
|---|---|
| **Dissociation reagent** | Accutase |

**Subculturing**

Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

**Thawing and culturing**

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

## QUALITY CONTROL AND CRYOPRESERVATION

**Sterility**

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

| | |
|---|---|
| **Freeze medium** | As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress. |

**Storage conditions**

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

**Shipping conditions**

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

## DONOR

| | |
|---|---|
| **Breed / subspecies** | C57BL/6 |

## LEGAL AND SAFETY

### Safety precautions

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

### Warranty

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

| | |
|---|---|
| **Material Transfer Agreement required** | Yes |