RAJI

300359

GENERAL INFORMATION

Description

Raji □胞是 R.J.V. Pulvertaft 于 1963 年从伯基特淋巴瘤中建立的淋巴母□胞样□胞系。这些□胞广泛应用于免疫学研究，因为它们高表达人类 CD19，CD19 可作为共受体，降低抗原 B □胞受体（BCR）刺激的阈值。Raji □胞不粘附，在悬浮液中以自由浮动的个体或双倍体形式生长。

这些□胞的倍增时间为 23.2 小时，直径相对较小，为 5-8 µm。Raji □胞的一些特点包括缺乏分化，因为它们会形成由数百个单个 □胞组成的大型聚集体。这些□胞是二倍体，核型稳定在雄性二倍体干系46内。

此外，Raji □胞对脊髓灰质炎病毒和水泡性口炎病毒有部分抵抗力。 Raji□胞高度表达人类CD19，已被确定为非霍奇金B□胞淋巴瘤中抗hCD19-CD3双特异性抗体的临床靶点。在 Raji 伯基特淋巴瘤□胞系和原发性淋巴瘤中也发现了 BCMA 的表达，使其成为免疫学家的一个重要研究领域。

Tissue	Maxilia
Disease	伯基特淋巴瘤

Synonyms

- Raji
- P1-Raji
- GM04671

CLASSIFICATION

Organism	Human
Sex	Male
Growth properties	Suspension
Biosafety level	1

CHARACTERISTICS

Cell type	淋巴母□胞

IDENTIFIERS

Citation

RAJI（Cytion 目录号 300359）

NCBI TaxID	9606
Cellosaurus	CVCL_0511

EXPRESSION AND SUSCEPTIBILITY

Products

在新城疫病毒的刺激下，□胞可能会产生干扰素。

HANDLING AND CULTURE

Culture medium

RPMI 1640，w：2.0 mM 稳定谷氨酰胺，w：2.0 g/L NaHCO3（Cytion 文章编号 820700a）。

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Incubation atmosphere	37°C, 5%CO2, 加湿环境。

Supplements

在培养基中添加 10% 热灭活 FBS

Subculturing

通过上下吸取操作轻柔匀化烧瓶中的□胞悬液，随后取代表性样本测定每毫升□胞密度。用新鲜培养基将悬液稀释至 1×10^5□胞/毫升浓度，并将调整后的悬液分装至新烧瓶中进行后续培养。

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

使用基于 PCR 的检测方法和基于发光的支原体检测方法排除支原体污染。
为确保没有□菌、真菌或酵母菌污染，每天都要对□胞培养物进行目视检查。

Freeze medium	我们使用完全生长培养基（包括 FBS）+10% DMSO 或 CM-1（Cytion 商品目录编号 800100）作为冷冻保存培养基，以获得足够的解冻后存活率，CM-1（Cytion 商品目录编号 800100）含有优化的渗透保护剂和代谢稳定剂，可提高恢复能力并减少冷冻引起的应激。

Storage conditions

如需长期保存，可将样品瓶放入气相液氮中，温度约为 -150 至 -196 ℃。在-80 °C下保存只能作为转入液氮前的短暂过渡。

Shipping conditions

冷冻保存的□胞系用干冰装运，并用经过验证的隔热包装和足够的制冷剂在整个运输过程中保持约 -78 °C 的温度。收到货后，应立即检查容器，并立即将小瓶转移到适当的储藏室。

DONOR

Donor age	11 年
Ethnicity	非洲, 尼日利亚

LEGAL AND SAFETY

Safety precautions

计划将低温样品储存在液氮中以便将来解冻时，必须遵守严格的安全措施。在将冷冻样本移入或移出液氮罐时，必须穿戴适当的防护手套和防护服，并使用面罩或安全护目镜。这样做是为了降低在移出时因冷冻样品爆炸而造成伤害的风险，因为爆炸可能导致锋利的碎片飞溅。

Warranty

在Cytion，我们保证产品的质量和活力。
我们保证，从我们的工厂发货后 60 天内，我们的□胞在开始培养时都是存活的。
我们保证所有材料均符合分析证书和/或产品说明书中描述的规格。
有关详□信息，请参阅我们的 <a href="" /Who-we-are/Warranty/"" hreflang=""x-default"">保修声明。

Material Transfer Agreement required	Yes

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB