MH-S

300487

GENERAL INFORMATION

Description

MH-S [illegible] [illegible] [illegible] [illegible] MH-S [illegible] [illegible] [illegible] [illegible] [illegible] MH-S [illegible] [illegible] [illegible] [illegible] [illegible] [illegible] [illegible] [illegible] MH-S [illegible] [illegible] [illegible]

Tissue	[illegible]

CLASSIFICATION

Organism	Mouse
Sex	Male
Growth properties	Adherent and suspension
Biosafety level	1

CHARACTERISTICS

Cell type	[illegible]

IDENTIFIERS

Citation

MH-S ([illegible] Cytion 300487)

NCBI TaxID	10090
Cellosaurus	CVCL_3855

EXPRESSION AND SUSCEPTIBILITY

Protein expression

[illegible] 1 (IL-1)

Antigen expression

[illegible]11[illegible] ([illegible]-1), [illegible] II ([illegible]-[illegible]), [illegible]

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Viruses

ทรานส์ฟอร์แมนท์ : ไวรัสลิง (SV40)

HANDLING AND CULTURE

Culture medium

RPMI 1640, w: 2.0 mM กลูตามีน □สถียร , w: 2.0 กรัม /ลิตร NaHCO3 (หมาย □ลขบทความ Cytion 820700a)

Incubation atmosphere

37°C, 5%CO2, บรรยากาศชื้น

Supplements

□ติมอาหาร □ลี้ยง □ชื้อด้วย FBS 10%

Dissociation reagent

แอคคู □ทส

Subculturing

รวบรวม □ซล □◌́ □□□ลอยในหลอดขนาด 15 มิลลิลิตร และล □าง □ซลล์ □◌ี □ติดผน □งอย่าง □บามือด้วย PBS □□□ ไม่มีแคล □ซียมและแมกนี □ซียม (ใช้ 3-5 มิลลิลิตรสำหรับขวด □ T25 และ 5-10 มิลลิลิตรสำหรับขวด □ T75)นำ□ Accutase (1-2 มล . สำ□หรับขวด T25, 2.5 มล . สำ□หรับขวด T75) ใส่ให้ทั่วชั้น □ซลล์อย่างทั่วถึง ปล่อยให้ □ซลล์พักที่อุณหภูมิห้อง □ป็น □วลา 10 นาที หลังจากนั้นรวมและปั่น □หวี่ยงทั้ง □ซลล์แขวนลอยและ □ซลล์ที่ □กาะติด หลังจากปั่น □หวี่ยงแล้ว ให้นำ □ □ซลล์ที่ตกตะกอนมาแขวนลอยอย่างระมัดระวังและถ่าย □ซลล์แขวนลอยไปยังขวดใหม่ที่มีอาหาร □ลี้ยง □ซลล์สด

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.

Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.

For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.

Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.

Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.

Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.

Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.

Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

การปน □ปื้อนของไมโคพลาสมาถูกตัดออกโดยใช้ทั้งการทดสอบแบบ PCR และการตรวจหาไมโคพลาสมาโดยใช้วิธีการ □รืองแสง

□พื่อให้แน่ใจว่าไม่มีการปน □ปื้อนของแบคที □รีย □ชื้อรา หรือยีสต์ การ □พาะ □ลี้ยง □ซลล์จะถูกตรวจสอบด้วยสายตาทุกวัน

Freeze medium

ในฐานะที่ □ป็นสารอาหารสำหรับการแช่□แข็ง □ราใช้สารอาหารสำ□หรับการ □จริญ □ติบโตแบบครบถ้วน (รวมถึง FBS) + 10% DMSO □พื่อความมีชีวิตหลังการละลายที่ □พียงพอ หรือ CM-1 (หมาย □ลขแคตตาล็อก Cytion 800100) ซึ่งรวมถึงสารป้องกันออสโมติกและสารรักษา □สถียรภาพ □มตาบอลิซึมที่ได้รับการปรับให้ □หมาะสม □พื่อ □พิ่มการฟื้นตัวและลดความ □ครียดที่ □กิดจากการแช่แข็ง

Storage conditions

สำ□หรับการ □ก็บรักษาในระยะยาว ให้วางหลอดบรรจุในไนโตร □จน □หลวในสถานะไอที่อุณหภูมิประมาณ −150 ถึง −196 ˚ C การ □ก็บรักษาที่อุณหภูมิ −80 °C สามารถใช้ได้ □ฉพาะในระยะสั้นก่อนการย้ายไปยังไนโตร □จน □หลว □ท่านั้น

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

2 / 3

Shipping conditions

[illegible] −78 °C [illegible]

DONOR

Breed / subspecies	BALB/cJ
Donor age	7 [illegible]

LEGAL AND SAFETY

Safety precautions

[illegible]

Warranty

[illegible]Cytion [illegible]
[illegible]
60 [illegible]
[illegible]
[illegible] [illegible]/[illegible]
[illegible]
<a href=""/Who-we-are/Warranty/"" hreflang=""x-default"">[illegible].

Material Transfer Agreement required	Yes