A2780

300491

GENERAL INFORMATION

Description

A2780 □□□□

GMO status details

No genetic modification; wildtype ovarian endometrioid carcinoma; parental line for A2780/DDP cisplatin-resistant derivative

Biosafety level	1

CHARACTERISTICS

Morphology	ลักษณะคล้าย □ยื่อบุผิว
Cell type	□ซลล์ □ยื่อบุผิว

IDENTIFIERS

Citation

A2780 (หมาย □ลขแคตตาล็อก Cytion 300491)

NCBI TaxID	9606
Cellosaurus	[CVCL_0134]

HANDLING AND CULTURE

Culture medium

RPMI 1640, w: 2.0 mM กลูตามีน □สถียร , w: 2.0 กรัม /ลิตร NaHCO3 (หมาย □ลขบทความ Cytion 820700a)

Incubation atmosphere	37°C, 5%CO2, บรรยากาศชื้น

Supplements

□ติมอาหาร □ลี้ยง □ชื้อด้วย FBS 10%

Dissociation reagent	แอคคู □ทส

Subculturing

รวบรวม □ซล □◌́ □□□ลอยในหลอดขนาด 15 มิลลิลิตร และล □าง □ซลล์ □◌ี □ติดผน □งอย่าง □บามือด้วย PBS □□□ ไม่มีแคล □ซียมและแมกนี □ซียม (ใช้ 3-5 มิลลิลิตรสำหรับขวด □ T25 และ 5-10 มิลลิลิตรสำหรับขวด □ T75)นำ□ Accutase (1-2 มล . สำ□หรับขวด T25, 2.5 มล . สำ□หรับขวด T75) ใส่ให้ทั่วชั้น □ซลล์อย่างทั่วถึง ปล่อยให้ □ซลล์พักที่อุณหภูมิห้อง □ป็น □วลา 10 นาที หลังจากนั้นรวมและปั่น □หวี่ยงทั้ง □ซลล์แขวนลอยและ □ซลล์ที่ □กาะติด หลังจากปั่น □หวี่ยงแล้ว ให้นำ □ □ซลล์ที่ตกตะกอนมาแขวนลอยอย่างระมัดระวังและถ่าย □ซลล์แขวนลอยไปยังขวดใหม่ที่มีอาหาร □ลี้ยง □ซลล์สด

Split ratio	1 ถึง 5
Seeding density	1 ถึง 3 × 10^4 □ซลล์ /ตาราง □ซนติ □มตร
Fluid renewal	2 ถึง 3 ครั้งต่อสัปดาห์

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

2 / 3

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

□□□