VCaP

300631

GENERAL INFORMATION

Description

The VCaP (Vertebral-Cancer of the Prostate) cell line is an important model in the study of prostate cancer, derived from a vertebral metastasis of a human prostate carcinoma. It was established to provide a relevant in vitro model for researching the biology of prostate cancer and its metastatic process, particularly focusing on hormone-refractory stages of the disease. VCaP cells are known for expressing a high level of prostate-specific antigen (PSA) and androgen receptor (AR), making them highly relevant for studies on androgen receptor signaling pathways and resistance mechanisms to anti-androgen therapy.
VCaP cells are also utilized extensively in genetic studies, as they harbor the TMPRSS2-ERG gene fusion, a common chromosomal translocation found in approximately 50% of prostate cancer cases. This specific genetic alteration is significant because it is thought to play a crucial role in the progression of prostate cancer. The cells are thus an excellent tool for research aiming to understand the molecular underpinnings of prostate cancer and for the development of new therapeutic strategies targeting TMPRSS2-ERG and related pathways. Moreover, VCaP cells exhibit robust in vitro growth and can form tumors when xenografted in immunodeficient mice, providing a useful system for preclinical studies of new anticancer drugs.
Overall, the VCaP cell line serves as a vital resource for molecular and pharmacological studies, contributing significantly to the understanding of prostate cancer biology and the assessment of new therapeutic agents. Its characteristics, including hormone responsiveness, gene fusion expression, and metastatic origin, make it uniquely suitable for advanced prostate cancer research, particularly in areas related to androgen independence and metastatic disease progression.

Tissue	Prostate
Disease	Prostate carcinoma
Metastatic site	Bone, vertebra

Synonyms

- VCAP
- Vcap
- Vertebral Cancer of the Prostate

CLASSIFICATION

Organism	Human
Sex	Male
Growth properties	Adherent
Biosafety level	1

CHARACTERISTICS

Tumourigenic	Yes, in SCID mice
Doubling time	Slow growing cell line, doubling time 5-6 days.

IDENTIFIERS

Citation

VCaP (Cytion catalog number 300631)

NCBI TaxID	9606

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Cellosaurus	CVCL_2235

EXPRESSION AND SUSCEPTIBILITY

Antigen expression

P53 antigen, Cytokeratin-18, prostate-specific antigen, prostatic acid phosphatase, Rb protein

Viruses

Mouse xenotropic retrovirus Bxv-1

HANDLING AND CULTURE

Culture medium

DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO3 (Cytion article number 820400a)

Incubation atmosphere	37°C, 5% CO_2, humidified atmosphere.

Supplements

Supplement the medium with 10% FBS

Dissociation reagent	Accutase

Subculturing

Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Seeding density	4-8 x 10^4 cells/cm^2

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Storage conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Shipping conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

DONOR

Donor age	59 years
Ethnicity	European

LEGAL AND SAFETY

Safety precautions

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Warranty

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Material Transfer Agreement required	Yes