HEK293 suspension-adapted

300686

□□ □□

□□

HEK293 □□□ □□ □□□□ □□ □□ □□ 293(HEK293) □□□ □□□□, □□ □□□ □□ □□□ □□□□ □□□□□ □□□□□□□. □□□ □□□ □□□ □□□ □ □□ □□□ □□ □□ □□□ □□□□□. □ □□□ □□□ □□□ □□□□ □□□ □□□ □□□□ □□ □ □□ □□□ □□□□ □□ HEK293 □□□ □□ □□□ □□□□, □ □ □□□□ □□ □□□ □□□ □□□□ □□□ □□□□ □□□ □ □□□□.
□ □□□ □□□ □□ □ □□ □□□ □□ □□□ □□□□ □□□□□ □□□□ □□ □□□□ □ □□ □□□□ □□ □□□ □□□□. □□ □□□ □□□□ □□□□ □□□□ □ □ □□□ □□□□□ □□□ □□□ □□□ □□□□□ □ □□□□□. HEK293 □□□ □□ □□□□ □□□□□□□, □□□□□□, □□□ □□ □□□□(AAV) □ □□□ □□ □□ □□ □□□□ □□□□, □□ □□ □□ □ □□□ □□□□ □□□ □□□.
□□□□□ HEK293 □□□ □□ □□□□ □□ □□□ □ □□□ □□□□ □□□□ □□□ □□□, □□□ □□□□ □□ □□□ □□□ □ □□ □□□ □□□□ □□□□□. □□ □ □□□ □□□□ □□ □□ □□□ □□ □□□□ □□□ □□ □ □□□ □□□□ □□□ □ □□ □□ □□ □□ □ □□ □□□□ □□□□ □ □ □□□□□.

□□	□□

□□ □□

□□ □□□

□□

□□□	□□
□	□□
□□ □□	□□
□□□ □□ □□	GMO-S1

□□□ □□ □□(GMO) □□ □□ □□

GMO-S1: This suspension-adapted HEK293 cell line contains adenovirus 5-derived E1 sequences from the parental HEK293 line, supporting high proliferative and protein expression capacity. The modification is stably present in transformed embryonic kidney cells. This classification applies only within Germany and may differ elsewhere.

□□□□ □□	1

□□

□□□	□□□
□□ □□□	□□ □□□□□

□□□

□□

HEK293 □□□□ □□□(□□□□ □□□□ □□ 300686)

NCBI □□ □□□□	9606
□□□□□□	CVCL_0045

□□ □ □□□

□□□ □□□
□□□□□

□□□ □□
CEA □□, p53 □□

□□□□ □□□
□□□□□□□ 5 DNA□ □□ □□□□□□□ 5 DNA

□□ □ □□

□□
□□□ 293S(□□□□□, □□)

□□ □□	37°C, 5%CO2, □□ □□.

□□□□□□
□□□ □□ □□

□□ □□	□□ □□

□□□
□□ □□□ 37°C/5% CO_2 □□□□□ □ □□□ □ □□□□□ □□ □□ □□□□□□ 5 x 10^5 ~ 2-3 x 10^6 □□/ml□ □□ □□□ □□□□□□. □□ □□□ 2-3 x 10^6 □□/ml□ □□□□ □ □ □□ □□□□□□. □□□ □□□ □□□ □□ □□□ □□□□□ □□□□□□. □□ □□□ 1-2 x 10^6 □□/ml□ □□□□, 200xg□□ 5□□ □□□ □□□ □□□ □□□□ □□□□ □□□□□□. □□□□ □□□□ □□□ □□□ □□□ □ □□□ □□□□ □□□□ □□ □□ □□ □□□ □□□□□. 200xg□□ 5□□ □□□ □□□ □□□ □□□□ □□□□ □□□□. □□□ □□ □□ □□□ □□□ □□□□□ □ □□ □□□□□. □□ □□□□ 80% □□□□□ □□, □□ □□□ 5-10□□ □□/ml□ □□□□□. □□ □□□□ □□□ □□□□ □□□ □□□□□□. 1°C/□□ □□ □□□ □□□□ □□ CoolCell □□ □□ □□ □□ □□ □□□□ □□□□□□.

□□ □□	5 x 10^5 □□/mL
□□ □ □□	□□ □□□ 5 x 10^5 □□/ml□ □□□ □□□□, □□□ □□□ □□ □□ □□□ 2-3 x 10^6 □□/ml□ □□□□□ □. 37°C/5% CO_2 □□□□ □□ □□□ □□ 100-150 rpm□□ □□□□□□.

□□ □ □□

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 200 x g for 5 minutes, carefully discard the supernatant containing freezing medium.
Follow the procedure described under Post-Thaw Recovery

□□ □□ □ □□ □□

□□
□□□□□□□ □□□ PCR □□ □□□□ □□ □□ □□□□□□□ □□□□ □□ □□□□ □□□□□.
□□□□, □□□ □□ □□ □□□ □□□ □□□□ □□ □□ □□□ □□ □□ □□□ □□□□.

□□□ □□	□□ □□ □□□ □□ □ □□□ □□□□ □□ □□ □□ □□ + 10% DMSO□ □□□□□.

□□ □□
□□ □□□ □□□□ □□□□ □ -150°C□□ -196°C□ □□□ □□ □□□ □□□□. 80°C□□ □□□□ □□ □□ □□□ □□□ □ □□ □□ □□□□ □□□□□.

□□ □□

□□ □□ □□□□ □□ □ □ -78°C□ □□□ □ □□ □□□ □□□ □□□ □□□ □□ □□□ □□□□□□□ □□ □□□□□. □□ □□ □□□ □□□□ □□□□ □□ □□ □ □□ □□ □□□ □□□□.

□□□

□□□ □□	□□

□□ □ □□

□□ □□

□□ □□□ □□ □□ □□□ □□ □□□ □□□□ □□, □□□ □□ □□□ □□□□ □□□. □□□ □□ □□□ □□□ □□□□, □□ □□ □□□ □□ □□□ □□ □□ □□□ □□ □□□□ □□□□ □□□□ □□□. □□ □□ □ □□□□ □□□ □□□□ □ □□ □□□ □□□ □□ □□ □□□ □□□ □□ □□□□.

□□

□□□□□ □□□ □□□ □□□□ □□□□□.
□□□ □□ □□□□ □□□ □ 60□ □□ □□ □□ □ □□□ □□□ □ □□□ □□□□□.
□□□ □□ □□□ □□ □□□ □/□□ □□ □□□ □□□ □□□ □□□□ □□□□□
□□□ □□□ <a href=""/Who-we-are/Warranty/"" hreflang=""x-default"">□□□ □ □□□□□ □□□□

□□ □□ □□□**(MTA)**□ □□□□□**.**	Yes