Ενδοθηλιακά κύτταρα της ανθρώπινης ομφάλιας αρτηρίας (HUAEC)

300698

ΓΕΝΙΚΈΣ ΠΛΗΡΟΦΟΡΊΕΣ

Περιγραφή

Primary Human Umbilical Artery Endothelial Cells (HUAEC) are isolated from the umbilical artery and provide an arterial counterpart to HUVEC. They are used to study arterial–venous differences in inflammation, nitric oxide signaling and fetal programming in pregnancy-related disorders.
Compared directly with HUVEC, HUAEC respond to inflammatory stimuli such as TNF-α with a distinct pattern of gene regulation, including differences in how they return to homeostasis after the stimulus is removed. This makes them well suited to test whether findings in venous endothelium also hold for arterial endothelium.

Ιστός	Umbilical cord (artery)
Ασθένεια	Normal umbilical artery endothelium (primary cells; non-tumorigenic; neonatal)
Θέση μετάστασης	Not applicable (normal, non-tumorigenic primary cell)

Εφαρμογές

Arterial–venous differences in inflammation and its resolution; Fetal programming: maternal obesity, oxidative stress and epigenetics; Nitric oxide, eNOS and arginase-2 regulation, epigenetic pharmacology; Angiogenesis assays (Matrigel tube formation, migration); Anti-proliferative screening of angiogenesis inhibitors (VEGF receptor 2)

Συνώνυμα

- HUAEC
- Human umbilical artery endothelial cells
- provitro 1210112

ΔΩΡΗΤΉΣ

Ηλικία δωρητή	Newborn (umbilical cord)
Εθνικότητα	Lot-specific (see CoA)

ΤΑΞΙΝΌΜΗΣΗ

Οργανισμός	Άνθρωπος
Σεξ	Σχετικά με το συγκεκριμένο παρτίδα
Ιδιότητες ανάπτυξης	Πιστός
Κατάσταση όσον αφορά τους ΓΤΟ	Χωρίς ΓΤΟ

Λεπτομέρειες σχετικά με την κατάσταση των ΓΤΟ

No genetic modification; primary human endothelial cells isolated from umbilical artery. Not transformed or immortalized. Tested and confirmed negative for mycoplasma, bacteria, yeast, HIV-1/2, HBV, HCV.

Επίπεδο βιοασφάλειας	1

ΧΑΡΑΚΤΗΡΙΣΤΙΚΆ

Μορφολογία	Cobblestone (endothelial)

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Τύπος κυττάρου	Primary endothelial cells
Ογκογόνος	No
Χρόνος διπλασιασμού	approx. 24 to 36 hours (lot-specific; guaranteed ≥15 population doublings under ScienCell-recommended conditions)

ΧΕΙΡΙΣΜΌΣ ΚΑΙ ΚΑΛΛΙΈΡΓΕΙΑ

Θρεπτικό μέσο

Endothelial Cell Growth Medium

Ατμόσφαιρα επώασης	37°C, 5% CO_2, saturated humidity.

Συμπληρώματα

ScienCell ECM #1001 is pre-supplemented with FBS, endothelial cell growth supplement (ECGS), and antibiotics as specified in the medium kit; no additional serum supplement needed. Coat culture vessels with fibronectin (1–2 µg/cm²) before seeding.

Αντιδραστήριο αποσύνδεσης	Subculture reagents as recommended on the certificate of analysis
Επίστρωση φιαλών	Fibronectin coating required (1–2 µg/cm²; incubate at 37°C for 30 min, aspirate excess before seeding)

Αναπαραγωγή σε υποκαλλιέργεια

Subculture the cells at about 75 % confluence. Denser cultures undergo irreversible contact inhibition and stop proliferating. Use the subculture reagents recommended on the certificate of analysis.

Αναλογία διαχωρισμού	1 to 3
Πυκνότητα σποράς	More than 6,000 cells/cm²
Ανανέωση υγρών	24 hours after seeding, then every 48 to 72 hours
Ανάκαμψη μετά την απόψυξη	Post-thaw viability ≥ 70 % viable cells. More than 80 % confluence within the expected recovery period.

Απόψυξη και καλλιέργεια

Thaw the cryovial quickly in a 37 °C water bath, swirling gently. Just before the last ice has melted, move the vial to a laminar flow hood and decontaminate its outer surface.
Transfer the cell suspension into a 15 ml centrifuge tube, add 10 ml of Endothelial Cell Growth Medium and centrifuge at 250 × g for 5 minutes.
Remove the supernatant and resuspend the cell pellet in 10 ml of Endothelial Cell Growth Medium. Check cell number and viability.
Divide the suspension between two 25 cm² culture flasks (5 ml each). Using two flasks reduces the risk of losing all cells if a passage fails.
Incubate at 37 °C, 5 % CO_2 and saturated humidity, with the screw caps opened by half a turn for gas exchange.
Change the medium after 24 hours to remove unattached cells, then every 48 to 72 hours, using pre-warmed medium only.
Subculture the cells at about 75 % confluence. Denser cultures undergo irreversible contact inhibition and stop proliferating. Use the subculture reagents recommended on the certificate of analysis.
Recommended seeding density: more than 6,000 cells/cm².

ΈΛΕΓΧΟΣ ΠΟΙΌΤΗΤΑΣ ΚΑΙ ΚΡΥΟΣΥΝΤΉΡΗΣΗ

Στειρότητα

Mycoplasma contamination is excluded by PCR.
To ensure there is no bacterial, fungal or yeast contamination, cultures are subjected to visual inspection.

Μέσο κατάψυξης	As a cryopreservation medium, we use complete growth medium + 10% DMSO for adequate post-thaw viability.

Συνθήκες αποθήκευσης

For long-term preservation, store vials in liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable for up to one week only.

Όροι αποστολής

Cryopreserved primary cells are shipped on dry ice. On receipt, check the cryovial for damage and transfer it without delay to appropriate storage.

ΈΚΦΡΑΣΗ ΚΑΙ ΕΥΑΙΣΘΗΣΊΑ

Επιφανειακά αντιγόνα

CD31 (PECAM-1) +; vWF (Factor VIII) +; Dil-Ac-LDL uptake +; Ephrin-B2 + (arterial marker); DLL4 + (arterial marker); Notch4 + (arterial marker); NRP1 + (arterial marker)

Υποδοχείς που εκφράζονται

VEGFR2 (KDR); VEGFR1 (Flt1); TIE-2; NRP1 (arterial-specific); Notch receptor (Notch1, Notch4)

Ιοί

Free of HIV-1, HIV-2, HBV, HCV (confirmed negative; ScienCell #8010 QC)

ΑΝΑΓΝΩΡΙΣΤΙΚΆ

Παραπομπή

Human umbilical artery endothelial cells (HUAEC) (Cytion catalog number 300698)

Αριθμός φορολογικού μητρώου NCBI	9606
Κελλοσαύρος	Not assigned

ΝΟΜΙΚΆ ΘΈΜΑΤΑ ΚΑΙ ΑΣΦΆΛΕΙΑ

Μέτρα ασφαλείας

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments. Primary cells are of human origin, and no known test procedure can ensure the complete absence of infectious agents. Handle all products as potentially infectious.

Εγγύηση

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Απαιτείται συμφωνία μεταφοράς υλικού	Yes