Sel otot polos arteri paru manusia (HPASMC)

300706

INFORMASI UMUM

Deskripsi

Primary Human Pulmonary Artery Smooth Muscle Cells (HPASMC) are isolated from the pulmonary artery. They are a key model for pulmonary arterial hypertension, including hypoxia-driven proliferation and vascular remodeling.
Hypoxia drives the proliferation of HPASMC, and PDGF-BB shifts them toward a synthetic, migratory phenotype, reproducing key features of pulmonary vascular remodeling. Because the balance of BMP and TGF-β signaling is central to pulmonary arterial hypertension, the cells are also used to study BMPR2-related mechanisms.

Tisu	Lung (pulmonary artery
Penyakit	Normal pulmonary arterial smooth muscle (non-tumorigenic primary cell)
Lokasi metastasis	Not applicable (normal, non-tumorigenic primary cell)

Aplikasi

Pulmonary arterial hypertension: BMP/TGF-β balance, proliferation and migration; Hypoxia-induced proliferation, PPARγ and microRNA-21; Long non-coding RNA regulation of migration in pulmonary arterial hypertension; Functional testing of BMPR2 promoter variants in reporter assays; PDGF-BB-induced phenotype switching and metabolomics; Endothelin-1 signaling and proteomics of BMPR2-mutant cells

Sinonim

- HPASMC
- Human pulmonary artery smooth muscle cells
- provitro 1210613

DONATUR

Usia donor	Lot-specific (see CoA)
Etnis	Lot-specific (see CoA)

KLASIFIKASI

Organisme	Manusia
Seks	Khusus lot
Sifat-sifat pertumbuhan	Pengikut
Status GMO	Bebas Transgenik

Rincian status GMO

No genetic modification; primary human pulmonary artery smooth muscle cells isolated from normal pulmonary arteries. Not transformed or immortalized.

Tingkat keamanan hayati	1

KARAKTERISTIK

Morfologi	Spindle-shaped (smooth muscle)
Jenis sel	Primary smooth muscle cells

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Tumorigenik	No
Waktu penggandaan	Not specified by AddexBio; typical primary VSMC doubling time 36–72 hours

PENANGANAN DAN BUDIDAYA

Media kultur

Smooth Muscle Cell Growth Medium kit (Cytion article number 850075); Smooth Muscle Cell Growth Medium mix (Cytion article number 850074)

Suasana inkubasi	37°C, 5% CO_2, saturated humidity.

Suplemen

Supplement as specified in the AddexBio SMC supplement kit (refer to specification sheet). For cryopreservation: complete culture medium + 5% (v/v) DMSO.

Reagen disosiasi	Subculture reagents as recommended on the certificate of analysis

Perbanyakan sel

Subculture the cells at about 75 % confluence. Denser cultures undergo irreversible contact inhibition and stop proliferating. Use the subculture reagents recommended on the certificate of analysis.

Rasio pemisahan	1 to 3
Kepadatan penanaman	More than 4,000 cells/cm²
Pergantian cairan	24 hours after seeding, then every 48 to 72 hours
Pemulihan pasca-pencairan	Post-thaw viability ≥ 70 % viable cells. More than 80 % confluence within the expected recovery period.

Pencairan dan pembiakan

Thaw the cryovial quickly in a 37 °C water bath, swirling gently. Just before the last ice has melted, move the vial to a laminar flow hood and decontaminate its outer surface.
Transfer the cell suspension into a 15 ml centrifuge tube, add 10 ml of Smooth Muscle Cell Growth Medium and centrifuge at 250 × g for 5 minutes.
Remove the supernatant and resuspend the cell pellet in 10 ml of Smooth Muscle Cell Growth Medium. Check cell number and viability.
Divide the suspension between two 25 cm² culture flasks (5 ml each). Using two flasks reduces the risk of losing all cells if a passage fails.
Incubate at 37 °C, 5 % CO_2 and saturated humidity, with the screw caps opened by half a turn for gas exchange.
Change the medium after 24 hours to remove unattached cells, then every 48 to 72 hours, using pre-warmed medium only.
Subculture the cells at about 75 % confluence. Denser cultures undergo irreversible contact inhibition and stop proliferating. Use the subculture reagents recommended on the certificate of analysis.
Recommended seeding density: more than 4,000 cells/cm².

PENGENDALIAN MUTU DAN KRIOPRESERVASI

Kemandulan

Mycoplasma contamination is excluded by PCR.
To ensure there is no bacterial, fungal or yeast contamination, cultures are subjected to visual inspection.

Media pembekuan	As a cryopreservation medium, we use complete growth medium + 10% DMSO for adequate post-thaw viability.

Kondisi penyimpanan

For long-term preservation, store vials in liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable for up to one week only.

Ketentuan pengiriman

Cryopreserved primary cells are shipped on dry ice. On receipt, check the cryovial for damage and transfer it without delay to appropriate storage.

EKSPRESI DAN KERENTANAN

Ekspresi protein

Smooth muscle α-actin; calponin; smooth muscle myosin heavy chain

Virus

Free of HIV, Hepatitis B, Hepatitis C (confirmed negative by AddexBio)

PENGIDENTIFIKASI

Referensi

Human pulmonary artery smooth muscle cells (HPASMC) (Cytion catalog number 300706)

Nomor Identifikasi Pajak NCBI	9606
Cellosaurus	Not assigned

HUKUM DAN KESELAMATAN

Tindakan pencegahan keselamatan

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments. Primary cells are of human origin, and no known test procedure can ensure the complete absence of infectious agents. Handle all products as potentially infectious.

Garansi

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Diperlukan Perjanjian Transfer Material	Yes

DETAIL PRODUK

Tabel komposisi