Cilvēka plaušu artēriju gludo muskuļu šūnas (HPASMC)

300706

VISPĀRĪGA INFORMĀCIJA

Apraksts

Primary Human Pulmonary Artery Smooth Muscle Cells (HPASMC) are isolated from the pulmonary artery. They are a key model for pulmonary arterial hypertension, including hypoxia-driven proliferation and vascular remodeling.
Hypoxia drives the proliferation of HPASMC, and PDGF-BB shifts them toward a synthetic, migratory phenotype, reproducing key features of pulmonary vascular remodeling. Because the balance of BMP and TGF-β signaling is central to pulmonary arterial hypertension, the cells are also used to study BMPR2-related mechanisms.

Audums	Lung (pulmonary artery
Slimība	Normal pulmonary arterial smooth muscle (non-tumorigenic primary cell)
Metastāžu vieta	Not applicable (normal, non-tumorigenic primary cell)

Pieteikumi

Pulmonary arterial hypertension: BMP/TGF-β balance, proliferation and migration; Hypoxia-induced proliferation, PPARγ and microRNA-21; Long non-coding RNA regulation of migration in pulmonary arterial hypertension; Functional testing of BMPR2 promoter variants in reporter assays; PDGF-BB-induced phenotype switching and metabolomics; Endothelin-1 signaling and proteomics of BMPR2-mutant cells

Sinonīmi

- HPASMC
- Human pulmonary artery smooth muscle cells
- provitro 1210613

ZIEDOTĀJS

Donora vecums	Lot-specific (see CoA)
Etniskā piederība	Lot-specific (see CoA)

KLASIFIKĀCIJA

Organisms	Cilvēks
Sekss	Atkarībā no partijas
Augšanas īpašības	Atbalstītājs
ĢMO statuss	Bez ĢMO

Sīkāka informācija par ĢMO statusu

No genetic modification; primary human pulmonary artery smooth muscle cells isolated from normal pulmonary arteries. Not transformed or immortalized.

Bioloģiskās drošības līmenis	1

RAKSTURLIELUMI

Morfoloģija	Spindle-shaped (smooth muscle)
Šūnu tips	Primary smooth muscle cells

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Tumorogēns	No
Laiks, kas nepieciešams, lai skaits dubultojās	Not specified by AddexBio; typical primary VSMC doubling time 36–72 hours

APSTRĀDE UN AUDZĒŠANA

Barotne

Smooth Muscle Cell Growth Medium kit (Cytion article number 850075); Smooth Muscle Cell Growth Medium mix (Cytion article number 850074)

Inkubācijas vide	37°C, 5% CO_2, saturated humidity.

Papildinājumi

Supplement as specified in the AddexBio SMC supplement kit (refer to specification sheet). For cryopreservation: complete culture medium + 5% (v/v) DMSO.

Disociācijas reaģents	Subculture reagents as recommended on the certificate of analysis

Pārstādīšana

Subculture the cells at about 75 % confluence. Denser cultures undergo irreversible contact inhibition and stop proliferating. Use the subculture reagents recommended on the certificate of analysis.

Sadalījuma attiecība	1 to 3
Sēšanas blīvums	More than 4,000 cells/cm²
Šķidruma atjaunošana	24 hours after seeding, then every 48 to 72 hours
Atgūšanās pēc atkusņa	Post-thaw viability ≥ 70 % viable cells. More than 80 % confluence within the expected recovery period.

Atkausēšana un kultivēšana

Thaw the cryovial quickly in a 37 °C water bath, swirling gently. Just before the last ice has melted, move the vial to a laminar flow hood and decontaminate its outer surface.
Transfer the cell suspension into a 15 ml centrifuge tube, add 10 ml of Smooth Muscle Cell Growth Medium and centrifuge at 250 × g for 5 minutes.
Remove the supernatant and resuspend the cell pellet in 10 ml of Smooth Muscle Cell Growth Medium. Check cell number and viability.
Divide the suspension between two 25 cm² culture flasks (5 ml each). Using two flasks reduces the risk of losing all cells if a passage fails.
Incubate at 37 °C, 5 % CO_2 and saturated humidity, with the screw caps opened by half a turn for gas exchange.
Change the medium after 24 hours to remove unattached cells, then every 48 to 72 hours, using pre-warmed medium only.
Subculture the cells at about 75 % confluence. Denser cultures undergo irreversible contact inhibition and stop proliferating. Use the subculture reagents recommended on the certificate of analysis.
Recommended seeding density: more than 4,000 cells/cm².

KVALITĀTES KONTROLE UN KRIOKONSERVĀCIJA

Sterilitāte

Mycoplasma contamination is excluded by PCR.
To ensure there is no bacterial, fungal or yeast contamination, cultures are subjected to visual inspection.

Saldēšanas barotne	As a cryopreservation medium, we use complete growth medium + 10% DMSO for adequate post-thaw viability.

Uzglabāšanas nosacījumi

For long-term preservation, store vials in liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable for up to one week only.

Piegādes nosacījumi

Cryopreserved primary cells are shipped on dry ice. On receipt, check the cryovial for damage and transfer it without delay to appropriate storage.

PAZĪMES UN UZŅĒMĪBA

Proteīnu ekspresija

Smooth muscle α-actin; calponin; smooth muscle myosin heavy chain

Vīrusi

Free of HIV, Hepatitis B, Hepatitis C (confirmed negative by AddexBio)

IDENTIFIKATORI

Atsauce

Human pulmonary artery smooth muscle cells (HPASMC) (Cytion catalog number 300706)

NCBI nodokļu identifikācijas numurs	9606
Cellosaurus	Not assigned

JURIDISKIE JAUTĀJUMI UN DROŠĪBA

Drošības pasākumi

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments. Primary cells are of human origin, and no known test procedure can ensure the complete absence of infectious agents. Handle all products as potentially infectious.

Garantija

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Nepieciešams materiālu nodošanas līgums	Yes

INFORMĀCIJA PAR PRODUKTU

Sastāva tabula