Cellules musculaires lisses bronchiques humaines (HBSMC)

300708

INFORMATIONS GÉNÉRALES

Description

Primary Human Bronchial Smooth Muscle Cells (HBSMC) are isolated from the bronchi. They are used to study airway remodeling, contractility and inflammatory signaling in asthma.
Airway smooth muscle cells are contractile and also immunomodulatory: in response to IL-13 they proliferate, migrate and produce periostin, and they release IL-6 and IL-8 upon immune stimulation. These responses link inflammation to the structural airway remodeling seen in asthma.

Tissu	Lung (bronchus)
Maladie	Normal bronchial smooth muscle (non-tumorigenic; primary cells)
Site métastatique	Not applicable (normal primary cells, not a tumor sample)

Applications

Asthma: chemokine receptors, wound repair and calcium signaling (asthmatic and healthy donors); IL-13-induced periostin production in Th2-driven airway inflammation; IL-13-driven proliferation and migration, store-operated calcium entry (Orai1); Immunoglobulin A-induced IL-6/IL-8 release and phenotype switching; Airway remodeling and smooth muscle dysfunction as a therapeutic target

Synonymes

- HBSMC
- Human bronchial smooth muscle cells
- provitro 1210632

DONATEUR

Âge du donneur	Lot-specific (see CoA)
Origine ethnique	Lot-specific (see CoA)

CLASSIFICATION

Organisme	Humain
Sexe	Spécifique au lot
Propriétés de croissance	Adepte
Statut OGM	Sans OGM

Informations détaillées sur le statut OGM

No genetic modification; primary human cells. Handle as potentially biohazardous material from human origin despite negative viral testing (HIV-1, HBV, HCV).

Niveau de biosécurité	1

CARACTÉRISTIQUES

Morphologie	Spindle-shaped (smooth muscle)
Type de cellule	Primary smooth muscle cells

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Tumorigène	No
Temps de doublement	approx. 36 to 48 hours (under standard culture conditions)

MANIPULATION ET CULTURE

Milieu de culture

Smooth Muscle Cell Growth Medium kit (Cytion article number 850075); Smooth Muscle Cell Growth Medium mix (Cytion article number 850074)

Atmosphère d'incubation	37°C, 5% CO_2, saturated humidity.

Compléments alimentaires

Smooth Muscle Cell Growth Supplement (ScienceCell catalog #1152) and 2% FBS included in SMCM. Alternatively, DMEM supplemented with 10% FBS, 2 mM L-glutamine, and 0.12 IU/mL insulin has been used in published studies.

Réactif de dissociation	Subculture reagents as recommended on the certificate of analysis

Repiquage

Subculture the cells at about 75 % confluence. Denser cultures undergo irreversible contact inhibition and stop proliferating. Use the subculture reagents recommended on the certificate of analysis.

Rapport de fractionnement	1 to 3
Densité de semis	More than 4,000 cells/cm²
Renouvellement des fluides	24 hours after seeding, then every 48 to 72 hours
Récupération après le dégel	Post-thaw viability ≥ 70 % viable cells. More than 80 % confluence within the expected recovery period.

Décongélation et mise en culture

Thaw the cryovial quickly in a 37 °C water bath, swirling gently. Just before the last ice has melted, move the vial to a laminar flow hood and decontaminate its outer surface.
Transfer the cell suspension into a 15 ml centrifuge tube, add 10 ml of Smooth Muscle Cell Growth Medium and centrifuge at 250 × g for 5 minutes.
Remove the supernatant and resuspend the cell pellet in 10 ml of Smooth Muscle Cell Growth Medium. Check cell number and viability.
Divide the suspension between two 25 cm² culture flasks (5 ml each). Using two flasks reduces the risk of losing all cells if a passage fails.
Incubate at 37 °C, 5 % CO_2 and saturated humidity, with the screw caps opened by half a turn for gas exchange.
Change the medium after 24 hours to remove unattached cells, then every 48 to 72 hours, using pre-warmed medium only.
Subculture the cells at about 75 % confluence. Denser cultures undergo irreversible contact inhibition and stop proliferating. Use the subculture reagents recommended on the certificate of analysis.
Recommended seeding density: more than 4,000 cells/cm².

CONTRÔLE QUALITÉ ET CRYOCONSERVATION

Stérilité

Mycoplasma contamination is excluded by PCR.
To ensure there is no bacterial, fungal or yeast contamination, cultures are subjected to visual inspection.

Milieu de congélation	As a cryopreservation medium, we use complete growth medium + 10% DMSO for adequate post-thaw viability.

Conditions de stockage

For long-term preservation, store vials in liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable for up to one week only.

Conditions de livraison

Cryopreserved primary cells are shipped on dry ice. On receipt, check the cryovial for damage and transfer it without delay to appropriate storage.

EXPRESSION ET SENSIBILITÉ

Antigènes de surface

α-smooth muscle actin (α-SMA) +; Desmin +

Expression des protéines

Smooth muscle α-actin; Desmin; Smooth muscle myosin heavy chain; Calponin; SM22; h-Caldesmon

Virus

Confirmed negative for HIV-1, HBV, HCV, mycoplasma, bacteria, yeast, and fungi (ScienceCell QC testing)

IDENTIFIANTS

Référence

Human bronchial smooth muscle cells (HBSMC) (Cytion catalog number 300708)

Numéro d'identification fiscale NCBI	9606
Cellosaurus	Not assigned (primary cells; individual donor; no Cellosaurus entry for primary HBSMC)

ASPECTS JURIDIQUES ET SÉCURITÉ

Consignes de sécurité

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments. Primary cells are of human origin, and no known test procedure can ensure the complete absence of infectious agents. Handle all products as potentially infectious.

Garantie

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Un accord de transfert de matériel est requis	Yes