HK EGFP-LaminB1/H2B-mCherry

300919

GENERAL INFORMATION

Description

The HK EGFP-LaminB1/H2B-mCherry cell line is an Hela Kyoto derived in vitro model designed for real-time visualization of chromatin dynamics and nuclear architecture in living cells. This cell line expresses two fluorescent protein fusions: EGFP (enhanced green fluorescent protein) fused with Lamin B1, and mCherry (a red fluorescent protein) fused with histone H2B. The fusion of EGFP with Lamin B1 allows for the observation of the nuclear envelope and nuclear lamina, structures critical for maintaining the integrity and functionality of the nucleus. Lamin proteins are type V intermediate filament proteins that form a meshwork underlying the inner nuclear membrane, playing key roles in nuclear stability, chromatin organization, and gene regulation.
On the other hand, the mCherry-tagged histone H2B enables the visualization of chromatin within the nucleus. Histones are fundamental components of the nucleosome, involved in the organization of DNA into chromatin, making them crucial for DNA replication, repair, and transcription. The mCherry tag on H2B provides a vivid red fluorescence that contrasts with the green fluorescence of EGFP, allowing for simultaneous dual-imaging of the nuclear structure and chromatin in live-cell experiments. This cell line is commonly used in studies focusing on nuclear mechanics, mitosis, and genome stability, providing a dynamic view of cellular processes that are otherwise difficult to observe in real time.

Tissue	Cervix
Disease	Carcinoma
Metastatic site	Primary tumor site (cervix)

Applications

Nuclear lamina and chromatin organization; Lamin B1 dynamics; H2B chromatin imaging; dual-color live-cell fluorescence; nuclear mechanics; mitosis; genome stability; nuclear envelope biology

Synonyms

- HeLa Kyoto EGFP-LaminB1 and H2B-mCherry

CLASSIFICATION

Organism	Human
Sex	Female
Growth properties	Adherent
GMO status	GMO-S1

GMO status details

GMO-S1: This HeLa Kyoto line contains EGFP-Lamin B1 and H2B-mCherry constructs for imaging nuclear envelope and chromatin organization. This classification applies only within Germany and may differ elsewhere.

Biosafety level	1

CHARACTERISTICS

Morphology	Epithelial-like cells with mosaic stone shape
Cell type	Epithelial cells

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

IDENTIFIERS

Citation

HK EGFP-LaminB1/H2B-mCherry (Cytion catalog number 300919)

NCBI TaxID	9606
Cellosaurus	CVCL_UR41
Depositor	The Ellenberg Lab (EMBL)

EXPRESSION AND SUSCEPTIBILITY

Protein expression

EGFP-LaminB1/H2B-mCherry

Products

Histone H2B

HANDLING AND CULTURE

Culture medium

DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO3, w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)

Incubation atmosphere	37°C, 5% CO_2, humidified atmosphere.

Supplements

Supplement the medium with 10% FBS

Dissociation reagent	Accutase

Subculturing

Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Seeding density	1 x 10^4 cells/cm^2
Fluid renewal	2 to 3 times per week
Post-thaw recovery	After thawing, plate the cells at 5 x 10^4 cells/cm^2 and allow the cells to recover from the freezing process and to adhere for at least 24 hours.

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Storage conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Shipping conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

DONOR

Donor age	30 years
Ethnicity	African American

LEGAL AND SAFETY

Safety precautions

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Warranty

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Third-party agreements apply	Yes