HK EGFP-LaminA/H2B-mCherry

300921

GENERAL INFORMATION

Description

HK EGFP-LaminA/H2B-mCherry □胞系是一种基因工程改造的 Hela Kyoto 衍生□胞模型，旨在促进活□胞内核动力学和染色质组织的高级研究。该 □胞系表达两种融合蛋白：EGFP（增强型绿色荧光蛋白）与层粘连蛋白 A 融合，mCherry（红色荧光蛋白）与组蛋白 H2B 融合。EGFP-Lamin A融合蛋白能突出核包膜，可观察 □胞周期进展过程中或各种实验条件下核结构的变化。与此同时，H2B-mCherry 融合蛋白与 DNA 结合，发出鲜艳的红色荧光，标记染色质，从而能够实时观察有丝分裂和间期的染色体过程。
这些□胞对于实时成像应用非常宝贵，包括研究核完整性、 DNA 复制和□胞衰老，以及研究核结构被破坏的疾病，如癌症和板层病。这种 □胞系的双色荧光功能允许同时观察核膜和染色质，有助于全面了解核-胞质相互作用和染色质的时空组织。这种功能使其成为分子生物学研究和 □胞生物物理学的重要工具，为基因表达调控机制、核组织和 □胞周期提供了洞察力。

Tissue	子宫颈
Disease	癌症

Synonyms

- HeLa Kyoto EGFP-LaminA and H2B-mCherry

CLASSIFICATION

Organism	Human
Sex	Female
Growth properties	Adherent
GMO status	GMO-S1

GMO status details

GMO-S1: This HeLa Kyoto line contains EGFP-Lamin A and H2B-mCherry constructs enabling dual-color imaging of nuclear lamina and chromatin. This classification applies only within Germany and may differ elsewhere.

Biosafety level	1

CHARACTERISTICS

Morphology	具有镶嵌石形状的上皮样□胞

IDENTIFIERS

Citation

HK EGFP-LaminA/H2B-mCherry（Cytion 目录号 300921）

NCBI TaxID	9606
Cellosaurus	CVCL_1D62
Depositor	艾伦伯格实验室（EMBL）

EXPRESSION AND SUSCEPTIBILITY

Protein expression

EGFP-LaminA/H2B-mCherry

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Products

组蛋白 H2B

HANDLING AND CULTURE

Culture medium

DMEM，w：4.5 克/升葡萄糖，w：4 毫摩尔 L-谷氨酰胺，w：3.7 克/升 NaHCO3，w：1.0 毫摩尔丙酮酸钠（Cytion 编号 820300a）。

Incubation atmosphere	37°C, 5%CO2, 加湿环境。

Supplements

在培养基中添加 10% FBS

Dissociation reagent	Accutase

Subculturing

去除粘附□胞上的旧培养基，用不含钙和镁的 PBS 冲洗。T25 烧瓶用 3-5 毫升 PBS，T75 烧瓶用 5-10 毫升。然后用 Accutase 完全覆盖□胞，T25 烧瓶用 1-2 毫升，T75 烧瓶用 2.5 毫升。让□胞在室温下孵育 8-10 分钟，使其分离。孵育后，用 10 毫升培养基轻轻混合□胞使其重悬，然后用 300xg 离心 3 分钟。弃去上清液，用新鲜培养基重悬 □胞，并将其转移到已装有新鲜培养基的新烧瓶中。

Seeding density	1 × 10^4 个□胞/平方厘米
Fluid renewal	每周 2 至 3 次
Post-thaw recovery	解冻后，将□胞以5×10^4个□胞/平方厘米的密度接种于培养皿中，使 □胞从冷冻过程中恢复并至少贴壁24小时。

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

使用基于 PCR 的检测方法和基于发光的支原体检测方法排除支原体污染。
为确保没有□菌、真菌或酵母菌污染，每天都要对 □胞培养物进行目视检查。

Freeze medium	我们使用完全生长培养基（包括 FBS）+10% DMSO 或 CM-1（Cytion 商品目录编号 800100）作为冷冻保存培养基，以获得足够的解冻后存活率，CM-1（Cytion 商品目录编号 800100）含有优化的渗透保护剂和代谢稳定剂，可提高恢复能力并减少冷冻引起的应激。

Storage conditions

如需长期保存，可将样品瓶放入气相液氮中，温度约为 -150 至 -196 ℃。在-80 °C下保存只能作为转入液氮前的短暂过渡。

Shipping conditions

冷冻保存的□胞系用干冰装运，并用经过验证的隔热包装和足够的制冷剂在整个运输过程中保持约 -78 °C 的温度。收到货后，应立即检查容器，并立即将小瓶转移到适当的储藏室。

DONOR

Donor age	30年

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Ethnicity	非洲裔美国人

LEGAL AND SAFETY

Safety precautions

计划将低温样品储存在液氮中以便将来解冻时，必须遵守严格的安全措施。在将冷冻样本移入或移出液氮罐时，必须穿戴适当的防护手套和防护服，并使用面罩或安全护目镜。这样做是为了降低在移出时因冷冻样品爆炸而造成伤害的风险，因为爆炸可能导致锋利的碎片飞溅。

Warranty

在Cytion，我们保证产品的质量和活力。
我们保证，从我们的工厂发货后 60 天内，我们的□胞在开始培养时都是存活的。
我们保证所有材料均符合分析证书和/或产品说明书中描述的规格。
有关详□信息，请参阅我们的 <a href="" /Who-we-are/Warranty/"" hreflang=""x-default"">保修声明。

Third-party agreements apply	Yes