B-LCL-CDG2

302013

GENERAL INFORMATION

Description

B-LCL-CDG2 □□□□□□□□□□□□□□□□□□□□□□□□□ B □□□□□□□□□□□□□□□□□□□□□□□ EBV □□□□□□□□□□□□□□□□□□□□□□□□□□□□□ □□□□□□ PMM2-CDG PMM2-CDG □□□ □□□□□□□□□□□□□□ □□□/□□□□□□□□□□□□□□□ □□□ □□ 2 (PMM2) □□□□□□□□□□□□□□ □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ PMM2

Tissue	□□□□□□□□□□□□□□□
Disease	□□

Applications

□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ CDG □□□□□□□□□□□□□□□□□□□□□ □□□□□□□□□□□□□□□□□□□□ (□□□□□ □□□□□□□□□ □□□□□□□□□□□ B) □□□□□□□□□□□□□□□□□□□□□ □□□□□□□□□□□□□□□□□□□□□□□□□□□□□ □□□□□□□□□□□□□□□□□□□□□□□□□□□□□ □□□□□□ □□□□□□□□□□□□□□ HLA □□□ □□□□□□□□□□□□□□□□□□□□□□□□□□□

CLASSIFICATION

Organism	Human
Sex	Female
Growth properties	Suspension
Biosafety level	2

CHARACTERISTICS

Morphology	□□□□□□□□□
Cell type	□□□□□□□□□□□□□

IDENTIFIERS

Citation

□□-□□□□□□□□□-□□□□□□□2 (□□□□□□□□□□□□□□□□□□ Cytion 302013)

NCBI TaxID	9606
Cellosaurus	CVCL_A9Y1

EXPRESSION AND SUSCEPTIBILITY

Surface antigens

CD60a- (GD3), CD60c-(7-O-acetylated GD3), CD75s+ sialylated lactosaminyl Noligosaccharides), CD77- (Gb3, globotriaosylceramide)

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Antigen expression

CD10-, CD19+, CD20+, CD21+, CD22+, CD23+, CD24+, CD37+m CD38+, CD39+, CD40+, CD53+, CD71+, CD72(+), CD73+,CD74 (+), CD80+, CD81+, CD82+, CD83-, CD84-, CD85+, CD86+, MHC class I+, MHC class II+

Viruses

ทราน □ ์ฟอ □ ์แมน □ ์ : EBV

HANDLING AND CULTURE

Culture medium

RPMI 1640, w: 2.0 mM กลูตามีน □สถียร , w: 2.0 กรัม /ลิตร NaHCO3 (หมาย □ลขบทความ Cytion 820700a)

Incubation atmosphere	37°C, 5%CO2, บรรยากาศชื้น

Supplements

□ติมอาหาร □ลี้ยง □ชื้อด้วย FBS ที่ทำให้ไม่□ทำงานทางชีวภาพด้□วยความร้อน 10%

Subculturing

รักษาวัฒนธรรมโดยการ □ติมหรือ □ปลี่ยนสื่อกลาง □ป็นระยะๆ □ริ่มต้นวัฒนธรรมด้วยความหนาแน่น 2 x 10^5 □ซลล์ /มล . และรักษาความ □ข้มข้นของ □ซลล์ให้อยู่ในช่วง 1 x 10^5 ถึง 5 x 10^5 □ซลล์ /มล . □พื่อการ □จริญ □ติบโตที่ □หมาะสมที่สุด

Fluid renewal	□มื่อสีกลาง □ปลี่ยน □ป็นสี □หลือง
Post-thaw recovery	ระดับกลาง

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

การปน □ปื้อนของไมโคพลาสมาถูกตัดออกโดยใช้ทั้งการทดสอบแบบ PCR และการตรวจหาไมโคพลาสมาโดยใช้วิธีการ □รืองแสง □พื่อให้แน่ใจว่าไม่มีการปน □ปื้อนของแบคที □รีย □ชื้อรา หรือยีสต์ การ □พาะ □ลี้ยง □ซลล์จะถูกตรวจสอบด้วยสายตาทุกวัน

Freeze medium	ในฐานะที่ □ป็นสารอาหารสำหรับการแช่□แข็ง □ราใช้สารอาหารสำ□หรับการ □จริญ □ติบโตแบบครบถ้วน (รวมถึง FBS) + 10% DMSO □พื่อความมีชีวิตหลังการละลายที่ □พียงพอ หรือ CM-1 (หมาย □ลขแคตตาล็อก Cytion 800100) ซึ่งรวมถึงสารป้องกันออสโมติกและสารรักษา □สถียรภาพ □มตาบอลิซึมที่ได้รับการปรับให้ □หมาะสม □พื่อ □พิ่มการฟื้นตัวและลดความ □ครียดที่ □กิดจากการแช่แข็ง

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

2 / 3

Storage conditions

[illegible] −150 [illegible] −196 °C [illegible] −80 °C [illegible]

Shipping conditions

[illegible] −78 °C [illegible]

DONOR

Donor age	[illegible]
Ethnicity	[illegible]

LEGAL AND SAFETY

Safety precautions

[illegible]

Warranty

[illegible]Cytion [illegible]
[illegible]
60 [illegible]
[illegible]
[illegible] [illegible]/[illegible]
[illegible]
<a href=""/Who-we-are/Warranty/"" hreflang=""x-default"">[illegible].

Material Transfer Agreement required	Yes