NCI-H1650

305059

GENERAL INFORMATION

Description

The NCI-H1650 cell line is derived from a human non-small cell lung carcinoma (NSCLC), specifically adenocarcinoma, and is widely used in cancer research due to its distinctive genetic profile and relevance in drug testing. This cell line features mutations in key oncogenic and tumor suppressor pathways, including a deletion in the PTEN gene and an activating mutation in EGFR. These genetic alterations make NCI-H1650 a suitable model for studying mechanisms of tumorigenesis and therapeutic resistance in NSCLC, especially in the context of targeted therapies aimed at the EGFR signaling pathway.

The deletion of PTEN in NCI-H1650 results in the loss of phosphatase activity, which deregulates the PI3K/AKT signaling pathway, contributing to tumor progression and resistance to certain therapeutic agents. The activating EGFR mutation, commonly observed in lung adenocarcinoma, renders the cell line particularly sensitive to tyrosine kinase inhibitors like erlotinib. However, the co-occurrence of these genetic changes often necessitates combination therapies to overcome adaptive resistance mechanisms that involve compensatory signaling pathways, such as mTOR or MET.

In addition to its genetic and signaling characteristics, NCI-H1650 has been included in numerous studies examining somatic mutations, copy number variations, and epigenetic alterations in cancer cell lines. Its response to inhibitors of EGFR and PI3K pathways highlights its utility in preclinical drug discovery and personalized medicine strategies. This cell line serves as a representative model for investigating the interplay between oncogenic drivers and therapeutic vulnerabilities in lung adenocarcinoma.

Tissue	Lung
Disease	Minimally invasive lung adenocarcinoma
Metastatic site	Pleural effusion

Synonyms

- NCI-H1650
- H-1650
- H1650_CO
- NCIH1650

CLASSIFICATION

Organism	Human
Sex	Male
Growth properties	Adherent
Biosafety level	1

CHARACTERISTICS

Morphology	Epithelial

IDENTIFIERS

Citation

NCI-H1650 (Cytion catalog number 305059)

NCBI TaxID	9606

Cellosaurus	CVCL_1483

HANDLING AND CULTURE

Culture medium

RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO3 (Cytion article number 820700a)

Incubation atmosphere	37°C, 5% CO_2, humidified atmosphere.

Supplements

Supplement the medium with 10% FBS

Dissociation reagent	Accutase

Subculturing

Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Fluid renewal	2 to 3 times per week

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Storage conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Shipping conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

DONOR

Donor age	27 years
Ethnicity	European

LEGAL AND SAFETY

Safety precautions

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Warranty

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Material Transfer Agreement required	Yes