CHO-HER2 magas

305413H

ÁLTALÁNOS INFORMÁCIÓK

Leírás

Disclaimer: The prices displayed for cell lines are exclusively for academic/not-for-profit customers. For commercial entitites the price is approximately €6,250.
If you represent a commercial entity or are unsure which category applies, please
contact us.
The CHO-HER2-High cell line is a stable recombinant CHO (Chinese Hamster Ovary) cell line engineered to express the HER2 receptor at a high level, approximately 85,000 molecules per cell. This cell line was generated using an innovative landing pad technology that ensures the HER2 gene is integrated at a specific, pre-validated genomic locus, allowing for consistent and reliable expression. HER2, also known as ERBB2 or CD340, is a member of the epidermal growth factor receptor (EGFR) family and plays a crucial role in regulating cell growth and differentiation. It is well-known for its involvement in breast and ovarian cancers, where its overexpression is linked to increased tumor aggressiveness and poorer patient outcomes. HER2 is a key target for cancer therapies such as Trastuzumab (Herceptin) and Pertuzumab (Perjeta). This cell line is versatile, supporting both adherent and suspension culture conditions, with adherent cells exhibiting an epithelial-like morphology. The expression of CXCR7 in this cell line was confirmed using flow cytometry.

Szövet	Ovary
Betegség	Chinese hamster ovary, non-neoplastic; genetically engineered for HER2 (ErbB2/CD340) surface expression (high expression level)

Alkalmazások

Antibody screening; ADCC/CDC assays; HER2-targeted therapy development; breast/gastric cancer research; flow cytometry

Szinonimák

- CHO-HER2

OSZTÁLYOZÁS

Organizmus	kínai hörcsög
Szex	Nő
Növekedési tulajdonságok	Adherens és szuszpenziós
GMO-státusz	GMO-S1

A GMO-státuszra vonatkozó részletek

GMO-S1: This CHO cell line contains a construct enabling high-level expression of human HER2 for oncology and receptor-signaling studies. This classification applies only within Germany and may differ elsewhere.

Biológiai biztonsági szint	1

JELLEMZŐK

Morfológia	Epithelial-like
Sejttípus	Epithelial cells
Kettőzési idő	approx. 14-16 hours

AZONOSÍTÓK

Hivatkozás

CHO-HER2 High (Cytion catalog number 305413H)

NCBI adóazonosító szám	10029
Cellosaurus	CVCL_A8W6

KIFEJEZŐDÉS ÉS FOGÉKONYSÁG

Kifejeződő receptorok

HER2

KEZELÉS ÉS TENYÉSZTÉS

Táptalaj

For adherent cultures: DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO3 (Cytion article number 820400a)
For suspension cultures: CHO Growth Medium A (from InSCREENeX; InSCREENeX catalog number INS-ME-1039)

Inkubációs környezet	37°C, 5% CO_2, humidified atmosphere.

Kiegészítők

For adherent cultures: Supplement the medium with 5% FBS. Add Geneticin (G418-Sulfat) to achieve a final concentration of 0.5 mg/mL.

Diszociációs reagens	For adherent cultures: Trypsin-EDTA

Átültetés

For routine adherent cell culture: Aspirate the old culture medium from the adherent cells, and wash them with PBS to remove any remaining medium. After aspirating the PBS, add the appropriate volume of Trypsin/EDTA solution based on the culture vessel size (e.g., 1 ml for a T25 flask, 3 ml for a T75 flask) and incubate at room temperature or 37°C for 5-10 minutes, or until the cells detach. Monitor detachment under a microscope, and gently tap the vessel if necessary to release the cells. Once detached, add complete medium to inactivate the Trypsin/EDTA, gently resuspend the cells, and transfer an aliquot of the cell suspension into a new culture vessel containing fresh medium. Place the vessel in an incubator set to 37°C with 5% CO_2, and change the medium every 2-3 days.

Osztási arány	1 to 5
Ültetési sűrűség	2 to 5 x 10^4 cells/cm^2
Folyadékcsere	2 to 3 times per week
Az olvadás utáni helyreállás	After thawing, split the cells at a ratio of 1:2 to 1:3 in T25 flasks and allow the cells to recover from the freezing process and to adhere (for adherent cultures) for at least 24 hours.

Felolvasztás és tenyésztés

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

MINŐSÉGELLENŐRZÉS ÉS KRIOKONZERVÁLÁS

Meddőség

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Fagyasztóközeg	As a cryopreservation medium, we recommend CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Tárolási feltételek

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Szállítási feltételek

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

ADOMÁNYOZÓ

Az adományozó életkora	Adult

JOGI ÉS BIZTONSÁGI INFORMÁCIÓK

Biztonsági óvintézkedések

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Jogi nyilatkozat

Our cells are provided for in vitro laboratory research purposes exclusively and are not intended for clinical or diagnostic use, nor are they to be administered to humans or used for veterinary purposes. Users must adhere to all applicable guidelines and regulations for the handling and use of these cells in a research setting.

Garancia

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Harmadik felekkel kötött megállapodások érvényesek	Yes