HEK293-CXCR7

305421

GENERAL INFORMATION

Description

□□□□□□□□□□□□□□□□□□□□/□□□□□□□□□□□□□□□□□ 6,250 □□
□□□□□□□□□□□□□□□□□□□□□□□□
□□□□□
HEK293-CXCR7□□□□□□□□□□□□HEK293□□□□□□□□□□□□□□□□□CXCR7□□□□□□□□□□inscreenex□“□□□”□□□□□□□□CXCR7□□□□□
□□□□□□□□□□□□□□□□□□□□□□□□□□□ CXCR7□□□ACKR3□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□G□□□□□□□□GPCR□□
□□CXCR7□□□G□□□□□□□□□□□□□□□CXCL12□CXCL11□□□□□□□□□CXCR4□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□
□□□□□□□□□
□□□□□□□□□□□□□□□□□□□□□□□□□□□CXCR7□□□

Tissue	□□□□
Disease	□□/□□□□□□□□□□HEK293□□□

Applications

CXCR7/ACKR3□□□□□□□□□GPCR□□□□□□□□□□□□□□□□CXCL11/CXCL12□□□□□□□□□□□□□□□□□□

CLASSIFICATION

Organism	Human
Sex	Female
Growth properties	Adherent
GMO status	GMO-S1

GMO status details

GMO-S1: This HEK293 line contains a CXCR7 expression construct, enabling studies of chemokine receptor activity. This classification applies only within Germany and may differ elsewhere.

Biosafety level	1

CHARACTERISTICS

Morphology	□□□
Cell type	□□□□
Doubling time	□24-36□□

IDENTIFIERS

Citation

HEK293-CXCR7□Cytion □□□ 305421□

NCBI TaxID	9606
Cellosaurus	CVCL_6G22

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

EXPRESSION AND SUSCEPTIBILITY

Receptors expressed

CXCR7□ACKR3□

HANDLING AND CULTURE

Culture medium

RPMI 1640□w□2.0 mM □□□□□□□w□2.0 g/L $NaHCO_3$□Cytion □□□□ 820700a□□

Incubation atmosphere	37°C, 5%CO2, □□□□□

Supplements

□□□□□□□ 10% FBS□1 mM □□□□□10 mM HEPES □ 1% NEAA□□□□□□□G418-□□□□□□□□□□□□ 1 mg/mL□

Dissociation reagent	□□□□-EDTA

Subculturing

□□□□□□□□□□□□□□□□□□□□□□□□□□□ PBS □□□□□□□□□□□□□□□□ PBS □□□□□□□□□□□□□□□□□□□□□□/EDTA □□□□ T25 □□ 1 □□□T75 □□ 3 □□□□□□□ 37°C □□□□□□□□□5-10 □□/EDTA□□ □□□ 37°C□□□□□□□□ 5%□□□□□□□□ 2-3 □□□□□□□□□□

Split ratio	1 □ 5
Seeding density	2□4×10⁴□□□/□□□□
Fluid renewal	□□ 2 □ 3 □
Post-thaw recovery	□□□□□ 1:2 □ 1:3 □□□□□□□□□ T25 □□□□□□□□□□□□□□□□□□□□□□□ 24 □□□ □□ □□□□□□□□□□□□□□□

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

□□□□ PCR □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□
□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

Freeze medium	□□□□□□□□□□□□□□□□ FBS□+10% DMSO □ CM-1□Cytion □□□□□□ 800100□□□□□□□□□ □□□□□□□□□□□□□□□□□□□CM-1□Cytion □□□□□□ 800100□□□□□□□□□□□□□□□□□□□□□□□ □□□□□□□□□□□□□□□□□□□□

Storage conditions

□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ -150 □ -196 ℃□□-80 °C□□□□□□□□□□□□□□□□□□□□□□

Shipping conditions
冷冻保存的细胞系用干冰装运，并用经过验证的隔热包装和足够的制冷剂在整个运输过程中保持约 -78 °C 的温度。收到货后，应立即检查容器，并立即将小瓶转移到适当的储藏室。

DONOR

Donor age 胎儿

LEGAL AND SAFETY

Safety precautions
计划将低温样品储存在液氮中以便将来解冻时，必须遵守严格的安全措施。在将冷冻样本移入或移出液氮罐时，必须穿戴适当的防护手套和防护服，并使用面罩或安全护目镜。这样做是为了降低在移出时因冷冻样品爆炸而造成伤害的风险，因为爆炸可能导致锋利的碎片飞溅。

Warranty

在Cytion，我们保证产品的质量和活力。
我们保证，从我们的工厂发货后 60 天内，我们的细胞在开始培养时都是存活的。
我们保证所有材料均符合分析证书和/或产品说明书中描述的规格。

有关详细信息，请参阅我们的 <a href="" /Who-we-are/Warranty/"" hreflang=""x-default"">保修声明。

Third-party agreements apply Yes

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

3 / 3