HEK293-TACD2

305424

GENERAL INFORMATION

Description

□□□□□□□□□□□□□□□□□□□□/□□□□□□□□□□□□□□□□□ 6,250 □□
□□□□□□□□□□□□□□□□□□□□□□□□
□□□□□
HEK293-TACD2□□□□□□□□□□□□HEK293□□□□□□□□□□□□□□□□□□TACD2□□□□□□□10,000□□□□□□□□□□□inscreenex□“□□□”□□□
□□□□TACD2□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ TACD2□□□TROP2□GA733-1□□□□□□□□□□□□□□□□□□□□□□□□□□□□
□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□TACD2□□□
□□
□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ TACD2 □□□□□□□□□□□□□□□□□□□□□□□□□□□□□

Tissue	□□□□
Disease	□□/□□□□□□□□□□HEK293□□□

Applications

□□TROP2□TACD2/TACSTD2□□□□□□□□□□□□□□□ADC□□□□sacituzumab govitecan□□□□□□□□□□□□□□□□□□□□□□□□□□□ADCC/CDC□
□□□□□□□□

CLASSIFICATION

Organism	Human
Sex	Female
Growth properties	Adherent
GMO status	GMO-S1

GMO status details

GMO-S1: This HEK293 line contains a TACD2 expression construct for receptor-binding and functional analyses. This classification applies only within Germany and may differ elsewhere.

Biosafety level	1

CHARACTERISTICS

Morphology	□□□
Cell type	□□□□
Doubling time	□24-36□□

IDENTIFIERS

Citation

HEK293-TACD2 □Cytion □□□ 305424□

NCBI TaxID	9606
Cellosaurus	CVCL_6G29

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

EXPRESSION AND SUSCEPTIBILITY

Receptors expressed

TACD2□TROP2 □ GA733-1□

HANDLING AND CULTURE

Culture medium

RPMI 1640□w□2.0 mM □□□□□□□w□2.0 g/L NaHCO3□Cytion □□□□ 820700a□□

Incubation atmosphere	37°C, 5%CO2, □□□□□

Supplements

□□□□□□□ 10% FBS□1 mM □□□□□10 mM HEPES □ 1% NEAA□□□□□□□G418-□□□□□□□□□□□□□ 1 mg/mL□

Dissociation reagent	□□□□-EDTA

Subculturing

□□□□□□□□□□□□□□□□□□□□□□□□□ PBS □□□□□□□□□□□□□□□ PBS □□□□□□□□□□□□□□□□□□□/EDTA □□□□ T25 □□ 1 □□□T75 □□ 3 □□□□□□□□ 37°C □□□□□□□□□□5-10 □□□□□□□□□□□□□□□□□□□□

Shipping conditions

□□□