Z-138

305516

GENERAL INFORMATION

Description

Z-138 is a human mantle cell lymphoma (MCL) cell line derived from a patient who initially presented with chronic lymphocytic leukemia (CLL) that later transformed into an aggressive B-cell leukemia. Z-138 cells exhibit a mature B-cell phenotype, expressing markers such as CD19, CD20, and surface immunoglobulin, while also maintaining CD5 positivity, a characteristic feature of mantle cell lymphoma. These cells have been extensively studied for their response to therapeutic agents, particularly in the context of BCL-2 inhibition. Studies have shown that Z-138 cells are sensitive to venetoclax, a selective BCL-2 inhibitor, which induces apoptosis through mitochondrial pathway activation. However, mechanisms of resistance to venetoclax have been identified, including upregulation of the PI3K/AKT pathway, highlighting the potential for combination therapies targeting both BCL-2 and PI3K/AKT signaling.
Z-138 cells have been utilized in research examining cytokine-mediated cytotoxicity, such as interleukin-21 (IL-21)-induced apoptosis. While some MCL cell lines are responsive to IL-21 treatment, Z-138 has demonstrated resistance, likely due to intrinsic differences in apoptotic signaling pathways. This resistance pattern provides insights into the heterogeneity of MCL and underscores the need for personalized therapeutic approaches. Z-138 serves as an important preclinical model for studying MCL pathogenesis and testing novel therapeutic strategies.

Tissue	Bone marrow
Disease	Mantle cell lymphoma

Applications

Immunology

Synonyms

- Z138

CLASSIFICATION

Organism	Human
Sex	Male
Growth properties	Suspension
Biosafety level	1

CHARACTERISTICS

Morphology	Lymphoblast
Cell type	Lymphoblast
Doubling time	24 hours

IDENTIFIERS

Citation

Z-138 (Cytion catalog number 305516)

NCBI TaxID	9606
Cellosaurus	CVCL_B077

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

EXPRESSION AND SUSCEPTIBILITY

Antigen expression

CD3-, CD5-, CD10-, CD19+, CD20+, CD23+, FMC7-

Mutational profile

Mutation: TRAF2, Simple, p.Trp114Ter (c.341G>A), Unspecified

HANDLING AND CULTURE

Incubation atmosphere	37°C, 5% CO_2, humidified atmosphere.

Subculturing

Maintain cultures by periodically adding or replacing the medium. Initiate cultures with a density of 5 x 10^5 cells/ml and keep the cell concentration within the range of 3 x 10^5 to 1 x 10^6 cells/ml for optimal growth.

Fluid renewal	2 to 3 times per week

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Storage conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Shipping conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

DONOR

Donor age	70 years
Ethnicity	Caucasian

LEGAL AND SAFETY

Safety precautions

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Warranty

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Material Transfer Agreement required	Yes

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB