NPC/HK1

305531

GENERAL INFORMATION

Description

NPC/HK1□□□□ HK-1 □ HK1□□□□□□□ 58 □□□□□□□□□□□□□□□□□□□□□NPC□□□□□ NPC/HK1 □□□□□□□□□□□□□□□□□□□□ 28 ± 2 □□□NPC/HK1 □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□-□□□□□EBV□□□□□□□□□□□□□□□□□□□□□□□□ □□□□□□□□□□□□□□□□□ □ EBV □□□□ EBV □□□□□□□□ EBV □□□□□□□□□□□□□□□□□□□□□□□□□□□ EBV □□□□□□
NPC/HK1 □□□□□□□□□□□□□□-□□□□□□□□□□□□□EBV □□□□□ EBV □□□□□□□□□□□□□□□□□□□□EGFR □□□□□□□□□□□□□□□□□□□□□□□□□□ □□ □□ □□□□EBV□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

Tissue	□□□□
Disease	□□□

Applications

□□□□□□EBV □□□□□□-□□□□□EMT□□□□□□□□□EGFR □□

Synonyms

- HK1
- HK-1

CLASSIFICATION

Organism	Human
Sex	Male
Growth properties	Adherent
Biosafety level	1

CHARACTERISTICS

Morphology	□□□
Cell type	□□□□
Doubling time	28 ± 2 □□

IDENTIFIERS

Citation

NPC/HK1□Cytion □□ 305531□

NCBI TaxID	9606
Cellosaurus	CVCL_7084

HANDLING AND CULTURE

Culture medium

RPMI 1640□□ 2.0 mM □□□□□□□□ 2.0 g/L $NaHCO_3$□Cytion □□□□ 820700a□

Incubation atmosphere	37°C□5% CO_2□□□□□□

Supplements

□□□□□□□ 10% □□□□□□FBS□

Dissociation reagent	Accutase

Subculturing

□□□□□□□□□□□□□□PBS□□□□□Accutase□□□□□□□□□ 8–10 □□□□□□□□□□□□□□ 300×g □□ 3 □□□□□□□□□□□□□□□□□□□□□□□□

Split ratio	1 □ 5
Seeding density	1 □ 3 × 10^4 □□□/cm²
Fluid renewal	□ 2 □ 3 □

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 200 x g for 5 minutes, carefully discard the supernatant containing freezing medium.
Follow the procedure described under Post-Thaw Recovery

QUALITY CONTROL AND CRYOPRESERVATION

Freeze medium	□□□□□□□□□□□□□□□□□□□□□□□□□ 10% □□□□□□□DMSO□□□□□□□□□□□□□□□□□□□□□□

Storage conditions

□□□□□□□□□□□□□□□□□□ −150 □ −196 °C □□□□□□□□□□□□□□□□□□□□□□□□□□□□ −80 °C □□□□□□□□□□□□□□

Shipping conditions

□□ −78 °C□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

DONOR

Donor age	58□
Ethnicity	□□

LEGAL AND SAFETY

Safety precautions

□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ □□□

Warranty

□ Cytion□□□□□□□□□□□□□□□□□□□□
□□□□□□□□□□□□□□□
60 □□□□□□□□□□□□□□□□□□□□□□□
□□□□□□□□□□□□
□□□□□□□□□□□□□□□□□□□□□
□□□□□□□□
□□□□□□□□

Material Transfer Agreement required	Yes