GL-261-Luc

305662

GENERAL INFORMATION

Description

□□□□□ GL261-Luc □□□□□□

IDENTIFIERS

Citation

GL-261-Luc (หมาย □ลขแคตตาล็อก Cytion 305662)

NCBI TaxID	10090
Cellosaurus	CVCL_C9CB

EXPRESSION AND SUSCEPTIBILITY

Antigen expression

Luc2 (ไฟลั๊ก , โคดอน -ออปติไมซ์)

HANDLING AND CULTURE

Culture medium

DMEM, w: 4.5 กรัม /ลิตร กลูโคส , w: 4 มิลลิโมลาร์ แอล -กลูตามีน , w: 3.7 กรัม /ลิตร NaHCO3, w: 1.0 มิลลิโมลาร์ โซ □ดียมไพรู □วต (หมาย □ลขบทความ Cytion 820300a)

Incubation atmosphere	37°C, 5%CO2, บรรยากาศชื้น

Supplements

□ติมอาหาร □ลี้ยง □ชื้อด้วย FBS 10%

Dissociation reagent	แอคคู □ทส

Subculturing

นำ□อาหาร □ลี้ยง □ซลล์ □ก่าออกจาก □ซลล์ที่ □กะติดและล้างด้วย PBS ที่ไม่มีแคล □ซียมและแมกนี □ซียม □□□หรับขวด T25 ให้ใช้ PBS 3-5 มล . และ □□□หรับขวด T75 ให้ใช้ 5-10 มล . จาก □□□นปิด □ซลล์ให้มิดด้วย Accutase โดยใช้ 1-2 มล . สำ□หรับขวด T25 และ 2.5 มล . สำ□หรับขวด T75ปล่อยให้ □ซลล์บ่มที่อุณหภูมิห้อง □ป็น □วลา 8-10 นาที □พื่อให้ □ซลล์หลุดออกจากกัน หลังจากบ่มแล้ว ให้ผสม □ซลล์ □บา ๆ กับอาหาร □ลี้ยง □ซลล์ 10 มิลลิลิตร □พื่อทำให้ □ □ซลล์แขวนลอยอีกครั้ง จากนั้นปั่น □หวี่ยงที่ 300 □ท่าของแรงโน้มถ่วงโลก (300xg) □ป็น □วลา 3 นาที □ทส่วนใสที่ได้ทิ้งไป แล้วแขวนลอย □ซลล์ในอาหาร □□ี □ยง □ซลล์ใหม่ จากนั้นถ่าย □ซลล์ไปยังขวด □พาะ □□□□ยงใหม่ □ี □□ติมอาหาร □□□□ยง □ซลล์ใหม่ไว้แล้ว

Seeding density	1 ถึง 3 x 10⁴ □ซลล์ /□ซนติ □มตร ²
Fluid renewal	2 ถึง 3 ครั้งต่อสัปดาห์

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.

Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.

For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.

Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.

Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.

Centrifuge the mixture at 200 x g for 5 minutes, carefully discard the supernatant containing freezing medium.

Follow the procedure described under Post-Thaw Recovery

QUALITY CONTROL AND CRYOPRESERVATION

Freeze medium	ในฐานะที่ □ป็นสารอาหารสำหรับการแช่□แข็ง □ราใช้สารอาหาร □□□หร □บการ □จริญ □ติบโตแบบครบถ้วน + 10% DMSO □□□□อความมีชีวิตหลังการละลายที่ □พียงพอ

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

2 / 3

Storage conditions

□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ □□□□□□□□□□□□□□□□□□□□