Neuro-2a-Luc

305690

GENERAL INFORMATION

Description

Neuro-2a-Luc [illegible] Neuro-2a (N2a) [illegible] crest [illegible] Neuro-2a [illegible] Neuro-2a-Luc [illegible] in vitro [illegible] in vivo [illegible]

[illegible] Neuro-2a-Luc [illegible] Neuro-2a [illegible]

[illegible] Neuro-2a-Luc [illegible]

Tissue	[illegible]
Disease	[illegible]

Synonyms

- Neuro2A-Luc

CLASSIFICATION

Organism	Mouse
Sex	Male
Growth properties	Adherent
GMO status	GMO-S1

GMO status details

GMO-S1: This cell line contains a stably integrated firefly luciferase reporter cassette (Luc2, codon-optimized) introduced via replication-incompetent lentiviral transduction. The resulting polyclonal cell population was maintained under puromycin selection (1–5 µg/mL). S1 containment is required. This classification applies only within Germany and may differ elsewhere.

Biosafety level	1

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

CHARACTERISTICS

Cell type	[illegible]

IDENTIFIERS

Citation

Neuro-2a-Luc ([illegible] Cytion 305690)

NCBI TaxID	10090
Cellosaurus	CVCL_K046

EXPRESSION AND SUSCEPTIBILITY

Antigen expression

Luc2 ([illegible], [illegible]-[illegible])

Viruses

[illegible] ([illegible]): [illegible]

Virus resistance

[illegible] 1

Reverse transcriptase	[illegible]

Products

[illegible], [illegible]

HANDLING AND CULTURE

Culture medium

EMEM (MEM Eagle), w: 2 mM L-Glutamine, w: 2.2 g/L $NaHCO_3$, w: EBSS ([illegible] Cytion 820100a)

Incubation atmosphere	37°C, 5%CO_2, [illegible]

Supplements

[illegible] FBS 10% [illegible] NEAA 1%

Dissociation reagent	[illegible]

Subculturing

[illegible] PBS [illegible] [illegible] T25 [illegible] PBS 3-5 [illegible]. [illegible] T75 [illegible] 5-10 [illegible]. [illegible] Accutase [illegible] 1-2 [illegible]. [illegible] T25 [illegible] 2.5 [illegible]. [illegible] T75[illegible] 8-10 [illegible] 10 [illegible] 300 [illegible] (300xg) [illegible] 3 [illegible]

Seeding density	1 [illegible] 3×10^4 [illegible]/[illegible]
Fluid renewal	2 [illegible] 3 [illegible]

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 200 x g for 5 minutes, carefully discard the supernatant containing freezing medium.
Follow the procedure described under Post-Thaw Recovery

QUALITY CONTROL AND CRYOPRESERVATION

Freeze medium	[illegible] + 10% DMSO [illegible]

Storage conditions

[illegible] −150 [illegible] −196 °C [illegible] −80 °C [illegible]

Shipping conditions

[illegible] −78 °C [illegible]

LEGAL AND SAFETY

Safety precautions

[illegible]

Warranty

[illegible]Cytion [illegible]
[illegible]
60 [illegible]
[illegible]
[illegible]/[illegible]
[illegible]
<a href=""/Who-we-are/Warranty/"" hreflang=""x-default"">[illegible].

Material Transfer Agreement required	Yes