SW620-Luc

305704

GENERAL INFORMATION

Description

SW620-Luc is a bioluminescent derivative of the human SW620 metastatic colon adenocarcinoma cell line, engineered to stably express a firefly luciferase reporter gene. The parental SW620 cell line was derived from a lymph node metastasis of the same patient from whom SW480 was established, allowing paired primary-metastatic comparative studies of colorectal cancer biology. SW620 cells exhibit epithelial-like morphology and share the same key oncogenic mutations as SW480, including KRAS G12V, TP53 R273H and P309S, and APC truncation, but display more mesenchymal features and greater invasive potential, consistent with their metastatic origin. SW620 is widely used to study colorectal cancer metastasis, epithelial-to-mesenchymal transition (EMT), and therapeutic resistance mechanisms.
The stable luciferase integration in SW620-Luc enables sensitive, quantitative bioluminescence imaging (BLI) of tumor burden in xenograft and experimental metastasis models in immunocompromised hosts. The emitted signal correlates with viable tumor cell number, supporting longitudinal monitoring of tumor engraftment, growth kinetics, liver colonization, and lymph node dissemination. SW620-Luc is particularly valuable for studying the metastatic progression of colorectal cancer, evaluating anti-metastatic agents, and performing comparative in vivo studies with the matched primary SW480 line.
SW620-Luc retains the molecular profile and metastatic phenotype of the parental SW620 line, including KRAS and TP53 mutations. The luciferase modification substantially enhances experimental throughput and sensitivity for in vivo pharmacodynamic assessment of treatment efficacy. Researchers should verify luciferase activity, mutational profile, and growth kinetics under their specific experimental conditions prior to large-scale preclinical use.

Tissue	Metastatic
Disease	Colon adenocarcinoma
Metastatic site	Lymph node

Synonyms

- SW620
- SW 620
- SW.620

CLASSIFICATION

Organism	Human
Sex	Male
Growth properties	Adherent
GMO status	GMO-S1

GMO status details

GMO-S1: This cell line contains a stably integrated firefly luciferase reporter cassette (Luc2, codon-optimized) introduced via replication-incompetent lentiviral transduction. The resulting polyclonal cell population was maintained under puromycin selection (1–5 µg/mL). S1 containment is required. This classification applies only within Germany and may differ elsewhere.

Biosafety level	1

CHARACTERISTICS

Morphology	Epithelial-like
Tumourigenic	Yes, in athymic nude mice

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

IDENTIFIERS

Citation

SW620-Luc (Cytion catalog number 305704)

NCBI TaxID	9606
Cellosaurus	CVCL_J268

EXPRESSION AND SUSCEPTIBILITY

Antigen expression

Luc2 (firefly, codon-optimized)

Mutational profile

Mutation: p.Gln1338Ter, Homozygous; Mutation: p.Gly12Val, Homozygous; Mutation: p.Arg273His, Heterozygous; Mutation: p.Pro309Ser, Heterozygous

HANDLING AND CULTURE

Culture medium

DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO3, w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)

Incubation atmosphere	37°C, 5% CO_2, humidified atmosphere.

Supplements

Supplement the medium with 10% FBS

Dissociation reagent	Accutase

Subculturing

Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Split ratio	1 to 3
Seeding density	1 to 3 x 10^4 cells/cm^2
Fluid renewal	2 to 3 times per week

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 200 x g for 5 minutes, carefully discard the supernatant containing freezing medium.
Follow the procedure described under Post-Thaw Recovery

QUALITY CONTROL AND CRYOPRESERVATION

Freeze medium	As a cryopreservation medium, we use complete growth medium + 10% DMSO for adequate post-thaw viability.

Storage conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Shipping conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

DONOR

Donor age	51 years
Ethnicity	Caucasian

LEGAL AND SAFETY

Safety precautions

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Warranty

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Material Transfer Agreement required	Yes