AKR-2B

305739

GENERAL INFORMATION

Description

AKR□□□□□ED-L2-cyclin D1; p53-/-□□□C57BL/6□□□□□□□□□□□□□□□□□□ESCC□□□□□ □□□□□□D1□□□□□□□□-□□□□ED-L2□□□□ □□□□□□-□□□□□□□□□□□□p53□□□□□□□□□-□□□□□□□□□□□□AKR□□□□
AKR□□□□□□□□□□□□□□□□□□□□□□□□□C57BL/6□□□□□□□□□□□□□□□□□□ □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ □□
AKR□□□□□□Hiroshi Nakagawa□□□□□·K·□□□□□Anil K. Rustgi□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

Tissue	□□
Disease	□□□□□□□

Applications

□□

Synonyms

- AKR-ECa

CLASSIFICATION

Organism	Mouse
Sex	Unspecified
Growth properties	Adherent
GMO status	Non-GMO

GMO status details

No

Biosafety level	1

CHARACTERISTICS

Morphology	□□□□
Tumourigenic	□□□□□□□□□C57BL/6□□□□□□□□□

IDENTIFIERS

Citation

AKR□Cytion □□□□ 305739□

NCBI TaxID	10090

EXPRESSION AND SUSCEPTIBILITY

Oncogenes

□EBV ED-L2□□□□□□Cyclin D1□CCND1□□□□

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Mutational profile

p53□Trp53□□□□□□

HANDLING AND CULTURE

Culture medium

DMEM□□4.5 g/L□□□□□4 mM L-□□□□

Incubation atmosphere	37°C, 5%CO2, □□□□□

Supplements

10% □□□□

Subculturing

□□□□□□□□80%□□□□0.05%□□□□/EDTA□□□□□□□□□□1:5□1:10□

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 200 x g for 5 minutes, carefully discard the supernatant containing freezing medium.
Follow the procedure described under Post-Thaw Recovery

QUALITY CONTROL AND CRYOPRESERVATION

Freeze medium	□□□□□□□□□□□□□□□□□□□□□□□+10% DMSO□□□□□□□□□□□□□□□

Storage conditions

□□□□□□□□□□□□□□□□□□□□□□□□□□ -150 □ -196 ℃□□-80 °C□□□□□□□□□□□□□□□□□□□

Shipping conditions

□□□ -78 °C □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

LEGAL AND SAFETY

Safety precautions

□□□□□□□□□□□□□□□□□□□