ChaGO-k-1

305759

ÁLTALÁNOS INFORMÁCIÓK

Leírás

The ChaGo-K-1 cell line is a clonal derivative of the ChaGo human bronchogenic carcinoma line, which was originally established from a primary lung tumor. This subline was developed through single-cell cloning of the parent ChaGo population, specifically selected for its elevated secretion of the α-subunit of human chorionic gonadotropin (HCG-α), a placental glycoprotein hormone subunit that is not typically produced by lung epithelial cells. In detailed hormone profiling, ChaGo-K-1 was shown to secrete extremely high levels of free HCG-α (approximately 290 pmol/day/mg protein), while secreting negligible levels of intact HCG or the β-subunit. This indicates that the cell line ectopically expresses HCG-α in the absence of normal trophoblastic function, a feature characteristic of some tumors exhibiting hormonal dysregulation.
Cytogenetic analysis of ChaGo-K-1 revealed a hyperdiploid karyotype with a modal chromosome number between 51 and 54, indicating chromosomal instability, which is typical of many human cancer cell lines. Importantly, electron microscopy demonstrated no evidence of mycoplasma contamination or viral particles, supporting the cell line's integrity and suitability for research. The unique ability of ChaGo-K-1 to ectopically secrete HCG-α at levels exceeding even the eutopic secretion of intact HCG by placental tumor cells (e.g., JEG-3) suggests that the mechanisms regulating glycoprotein hormone subunit expression can be co-opted or dysregulated in non-placental malignancies. This property makes ChaGo-K-1 an important model for studying ectopic hormone production, oncogenic dedifferentiation, and tumor-associated glycoprotein signaling.

Szövet	Metastatic
Betegség	Bronchogenic carcinoma

Alkalmazások

3D cell culture, Cancer research

Szinonimák

- CHAGO-K-1
- ChaGo K-1
- ChaGo-K1
- CHAGOK1
- ChaGoK1

OSZTÁLYOZÁS

Organizmus	Ember
Szex	Férfi
Növekedési tulajdonságok	Hűséges
Biológiai biztonsági szint	1

AZONOSÍTÓK

Hivatkozás

ChaGO-k-1 (Cytion catalog number 305759)

NCBI adóazonosító szám	9606
Cellosaurus	CVCL_1121

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

KIFEJEZŐDÉS ÉS FOGÉKONYSÁG

Antigén-expresszió

HCG; human chorionic gonadotropin (alpha subunit only); estradiol; progesterone; mucin (apomucin, MUC-1, MUC-2)

Izoenzimek

AK-1, 1,ES-D, 1,G6PD, B,GLO-I, 1-2,LDH, 5,Me-2, 0,PGM1, 1,PGM3, 1

Mutációs profil

Mutation: ARID1A, Simple, c.1921-2A>G, Heterozygous, Note=Splice acceptor mutation (Cosmic-CLP=687596), ARID1A, Simple, p.His684Asp (c.2050C>G), Heterozygous (Cosmic-CLP=687596), RB1, Simple, p.Glu837Lys (c.2509G>A), Homozygous (Cosmic-CLP=687596), TP53, Simple, p.Cys275Phe (c.824G>T), Heterozygous (Cosmic-CLP=687596)

KEZELÉS ÉS TENYÉSZTÉS

Inkubációs környezet	37°C, 5% CO_2, humidified atmosphere.
Folyadékcsere	2 to 3 times per week

Felolvasztás és tenyésztés

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

MINŐSÉGELLENŐRZÉS ÉS KRIOKONZERVÁLÁS

Meddőség

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Fagyasztóközeg	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Tárolási feltételek

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Szállítási feltételek

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

ADOMÁNYOZÓ

Az adományozó életkora	45 years
Etnikai hovatartozás	Caucasian

JOGI ÉS BIZTONSÁGI INFORMÁCIÓK

Biztonsági óvintézkedések

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Jogi nyilatkozat

Our cells are provided for in vitro laboratory research purposes exclusively and are not intended for clinical or diagnostic use, nor are they to be administered to humans or used for veterinary purposes. Users must adhere to all applicable guidelines and regulations for the handling and use of these cells in a research setting.

Garancia

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Anyagátadási megállapodás szükséges	Yes

TERMÉKADATOK

Összetételi táblázat