NCI-H322

305839

GENERAL INFORMATION

Description

NCI-H322 is a human non-small cell lung cancer (NSCLC) cell line derived from an adult patient with a bronchioalveolar carcinoma, a histologic subtype of adenocarcinoma. This cell line was established by the NCI-Navy Medical Oncology Branch as part of a comprehensive effort to generate clinically annotated lung cancer models for research and therapeutic development. NCI-H322 exhibits adherent epithelial morphology in vitro and is typically maintained in RPMI-1640 medium supplemented with 10% fetal bovine serum under standard cell culture conditions.

Molecular profiling of NCI-H322 reveals that it carries a KRAS mutation, which contributes to oncogenic signaling through the MAPK/ERK and PI3K/AKT pathways. This mutation renders the cell line resistant to EGFR-targeted therapies and makes it suitable for studies focused on KRAS-driven lung adenocarcinoma. Additionally, the line is wild-type for EGFR and TP53, offering a defined genetic context for dissecting KRAS-dependent tumor biology. Its transcriptional and proteomic data have been included in large-scale datasets such as the Cancer Cell Line Encyclopedia (CCLE), where it has contributed to analyses of lineage-specific vulnerabilities and drug response patterns.

NCI-H322 has been used extensively in pharmacologic screening and mechanistic studies to explore sensitivity to MEK inhibitors, PI3K pathway inhibitors, and chemotherapeutic agents. Its consistent performance across studies and well-documented mutation profile make it a valuable preclinical model for KRAS-mutant NSCLC, as well as a key reference in efforts to understand tumor heterogeneity and drug resistance in lung adenocarcinoma.

Tissue	Lung
Disease	Minimally invasive lung adenocarcinoma

Synonyms

- H322
- H-322
- H322T
- NCI-H322T
- NCIH322T
- NCI-322
- NCIH322

CLASSIFICATION

Organism	Human
Sex	Male
Growth properties	Adherent
Biosafety level	1

CHARACTERISTICS

Cell type	Club cells
Doubling time	50

IDENTIFIERS

Citation

NCI-H322 (Cytion catalog number 305839)

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

NCBI TaxID	9606
Cellosaurus	CVCL_1556

EXPRESSION AND SUSCEPTIBILITY

Mutational profile

Mutation: TP53, Simple, p.Arg248Leu (c.743G>T), Homozygous (PubMed=1311061, PubMed=1565469, PubMed=10536175, PubMed=20557307).

HANDLING AND CULTURE

Culture medium

RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO3 (Cytion article number 820700a)

Incubation atmosphere	37°C, 5% CO_2, humidified atmosphere.

Supplements

Supplement the medium with 10% FBS

Dissociation reagent	Accutase

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Storage conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Shipping conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

DONOR

Donor age	52 years

Ethnicity	Caucasian

LEGAL AND SAFETY

Safety precautions

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Disclaimer

Our cells are provided for in vitro laboratory research purposes exclusively and are not intended for clinical or diagnostic use, nor are they to be administered to humans or used for veterinary purposes. Users must adhere to all applicable guidelines and regulations for the handling and use of these cells in a research setting.

Warranty

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

Material Transfer Agreement required	Yes

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB