B95-8

601102

GENERAL INFORMATION

Description

B95-8 □胞系是一种永生化的狨猴 B 淋巴母□胞系，来源于棉顶狨猴（Saguinus oedipus）的外周血白□胞。这种□胞系是通过感染爱泼斯坦-巴氏病毒（EBV）建立的，这是一种使 B □胞永生的常用方法。EBV 的存在是 B95-8 □胞系在研究中发挥作用的关键，尤其是在与病毒肿瘤学、病毒 -宿主相互作用以及 EBV 本身的生物学相关的研究中。

在病毒学研究中，B95-8 □胞经常被用作 Epstein-Barr 病毒的来源。它们能产生传染性病毒颗粒，是繁殖 EBV 和进行需要活性病毒的实验的宝贵工具。此外，这种 □胞系还有助于开发疫苗和治疗策略，以预防 EBV 相关疾病，包括伯基特淋巴瘤和霍奇金淋巴瘤。这种 □胞还与研究 EBV 的免疫反应有关，因为它们可以用来模拟 B □胞的转化，并了解 EBV 诱导肿瘤发生的机制。

Tissue	血液

Synonyms

- B95.8
- B 95.8
- B 95-8
- B-95-8
- B958
- GM07404
- GM07404A
- GM07404D

CLASSIFICATION

Organism	Cotton-top tamarin
Sex	Female
Growth properties	Suspension
Biosafety level	2

CHARACTERISTICS

Morphology	淋巴母□胞

IDENTIFIERS

Citation

B95-8（Cytion 产品编号 601102）

NCBI TaxID	9490
Cellosaurus	CVCL_1953

HANDLING AND CULTURE

Culture medium

RPMI 1640，w：2.0 mM 稳定谷氨酰胺，w：2.0 g/L NaHCO3（Cytion 文章编号 820700a）。

Incubation atmosphere	37°C, 5%CO2, 加湿环境。

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

Supplements

在培养基中添加 10% FBS

Subculturing

通过上下吸取操作轻柔匀化烧瓶中的□胞悬液，随后取代表性样本测定每毫升□胞密度。用新鲜培养基将悬液稀释至 1×10^5□胞/毫升浓度，并将调整后的悬液分装至新烧瓶中进行后续培养。

Fluid renewal	每周 2 至 3 次

Thawing and culturing

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

QUALITY CONTROL AND CRYOPRESERVATION

Sterility

使用基于 PCR 的检测方法和基于发光的支原体检测方法排除支原体污染。
为确保没有□菌、真菌或酵母菌污染，每天都要对□胞培养物进行目视检查。

Freeze medium	我们使用完全生长培养基（包括 FBS）+10% DMSO 或 CM-1（Cytion 商品目录编号 800100）作为冷冻保存培养基，以获得足够的解冻后存活率，CM-1（Cytion 商品目录编号 800100）含有优化的渗透保护剂和代谢稳定剂，可提高恢复能力并减少冷冻引起的应激。

Storage conditions

如需长期保存，可将样品瓶放入气相液氮中，温度约为 -150 至 -196 ℃。在-80 °C下保存只能作为转入液氮前的短暂过渡。

Shipping conditions

冷冻保存的□胞系用干冰装运，并用经过验证的隔热包装和足够的制冷剂在整个运输过程中保持约 -78 °C 的温度。收到货后，应立即检查容器，并立即将小瓶转移到适当的储藏室。

LEGAL AND SAFETY

Safety precautions

计划将低温样品储存在液氮中以便将来解冻时，必须遵守严格的安全措施。在将冷冻样本移入或移出液氮罐时，必须穿戴适当的防护手套和防护服，并使用面罩或安全护目镜。这样做是为了降低在移出时因冷冻样品爆炸而造成伤害的风险，因为爆炸可能导致锋利的碎片飞溅。

Disclaimer

我们提供的□胞仅用于体外实验室研究目的，不用于临床或诊断用途，也不得用于人类或兽医目的。用户必须遵守在研究环境中处理和使用这些 □胞的所有适用准则和法规。

Warranty

在Cytion，我们保证产品的质量和活力。
我们保证，从我们的工厂发货后 60 天内，我们的□胞在开始培养时都是存活的。
我们保证所有材料均符合分析证书和/或产品说明书中描述的规格。
有关详□信息，请参阅我们的 <a href="" /Who-we-are/Warranty/"" hreflang=""x-default"">保修声明。

Material Transfer Agreement required	Yes

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB