CV-1

605471

INFORMAZIONI GENERALI

Descrizione

CV-1 is an African green monkey cell line derived from the kidney in 1964. Initially used in research that focused on the transformation of the cancerogenic Rous sarcoma virus (RSV), this fibroblast-like cell line is widely used in biological research for virus production, transfection, and gene silencing.
These cells are negative for reverse transcriptase and being susceptible to several viruses, including poliovirus 1, herpes simplex, simian virus 40 (SV40), California encephalitis, and both Eastern and Western equine encephalitis.
The CV-1 cell line exhibits rapid growth, grows adherent on plastic and glass surfaces and shows chromosome number shifts at high passage levels. It has been observed that CV-1 cells exhibit increased tumorigenicity in Wistar rats treated with ATG as well as increased cell colony formation in soft agar.
Moreover, CV-1 cells support the replication of SV40 virus and exhibit rapid thymidine kinase (TK) activity following induction of simian, adeno, and papovavirus infections. The karyotype of CV-1 cells is 2n = 60, pseudodiploid.
CV-1 cells have been used in a variety of specific applications in biological research, including efficacy testing, transfection host, and viruscide testing. They are also known to be a suitable host for transfection, especially by SV40 vectors.

Tessuto	Kidney

Applicazioni

Suitable host for transfection, especially by SV40 vectors.

Sinonimi

- Cv-1
- CV 1
- CV-1.K
- CV1

CLASSIFICAZIONE

Organismo	Scimmia verde africana
Sesso	Maschio
Proprietà di crescita	Aderente
Livello di biosicurezza	1

CARATTERISTICHE

Tipo di cellula	Fibroblast

IDENTIFICATORI

Riferimento bibliografico

CV-1 (Cytion catalog number 605471)

Codice fiscale NCBI	9534
Cellosaurus	CVCL_0229

Cytion GmbH
Nikola-Tesla-Str. 3
69124 Heidelberg
Tel: +49 (0)6221 405780
info@cytion.com
cytion.com

Registered office Heidelberg
VAT ID: DE283292917
Local court Mannheim - HRB 714394
CEO / Geschäftsführer: Jonathan Steubing

Banking information
Sparkasse Heidelberg
IBAN: DE35 6725 0020 0009 2076 86
BIC: SOLADES1HDB

ESPRESSIONE E SUSCETTIBILITÀ

Suscettibilità ai virus

Poliovirus 1, herpes simplex, Eastern equine encephalitis, Western equine encephalitis, California encephalitis, SV40

Trascriptasi inversa	Negative

MANIPOLAZIONE E COLTURA

Terreno di coltura

EMEM (MEM Eagle), w: 2 mM L-Glutamine, w: 2.2 g/L NaHCO3, w: EBSS (Cytion article number 820100a)

Atmosfera di incubazione	37°C, 5% CO_2, humidified atmosphere.

Integratori

Supplement the medium with 10% FBS and 1% NEAA

Reagente di dissociazione	Accutase

Subcoltura

Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Densità di semina	3 to 4 x 10^4 cells/cm^2 will yield in a confluent layer in about 4 days
Rinnovo dei fluidi	2 times per week
Recupero post-scongelamento	After thawing, plate the cells at 5 x 10^4 cells/cm^2 and allow the cells to recover from the freezing process and to adhere for at least 24 hours.

Scongelamento e coltura

Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

CONTROLLO DI QUALITÀ E CRIOCONSERVAZIONE

Sterilità

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.
To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Terreno di coltura congelato	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Condizioni di conservazione

For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Condizioni di spedizione

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

DONATORE

Età dei donatori	141 days

ASPETTI LEGALI E DI SICUREZZA

Misure di sicurezza

When planning to store a cryovial in liquid nitrogen for future thawing, it is mandatory to adhere to stringent safety measures. Appropriate protective gloves and clothing are essential, and the use of a face mask or safety goggles is required during the transfer of frozen samples to or from the liquid nitrogen tank. This is to mitigate the risk of injury from potential cryovial explosions upon removal, which can result in the projection of sharp fragments.

Dichiarazione di non responsabilità

Our cells are provided for in vitro laboratory research purposes exclusively and are not intended for clinical or diagnostic use, nor are they to be administered to humans or used for veterinary purposes. Users must adhere to all applicable guidelines and regulations for the handling and use of these cells in a research setting.

Garanzia

At Cytion, we stand behind the quality and viability of our products.
We guarantee that our cells will be viable at the start of culture for
60 days after shipment from our facilities.
We assure that all materials conform to the specifications described in the
Certificate of Analysis and/or Product Sheet.
For full details, please refer to our
Warranty Statement.

È richiesto un accordo di trasferimento di materiale	Yes